

BIHEP-TH-94-20

June, 1994

q -Deformed Chern Characters for Quantum Groups $SU_q(N)$

Bo-Yu Hou¹⁾, Bo-Yuan Hou²⁾ and Zhong-Qi Ma³⁾

1) Institute of Modern Physics, Northwest University, Xi'an 710069, P. R. China

2) Graduate School, Chinese Academy of Sciences, P. O. Box 3908, Beijing 100039, P. R. China

3) Institute of High Energy Physics, P. O. Box 918(4), Beijing 100039, P. R. of China

Abstract

In this paper, we introduce an $N \times N$ matrix $\epsilon^{a\bar{b}}$ in the quantum groups $SU_q(N)$ to transform the conjugate representation into the standard form so that we are able to compute the explicit forms of the important quantities in the bicovariant differential calculus on $SU_q(N)$, such as the q -deformed structure constant $\mathbf{C}_{IJ}^K$ and the q -deformed transposition operator Λ . From the q -gauge covariant condition we define the generalized q -deformed Killing form and the m -th q -deformed Chern class P_m for the quantum groups $SU_q(N)$. Some useful relations of the generalized q -deformed Killing form are presented. In terms of the q -deformed homotopy operator we are able to compute the q -deformed Chern-Simons Q_{2m-1} by the condition $dQ_{2m-1} = P_m$. Furthermore, the q -deformed co-cycle hierarchy, the q -deformed gauge covariant Lagrangian, and the q -deformed Yang-Mills equation are derived.

1. INTRODUCTION

Recently, quantum groups have attracted increasing attention. Since the quantum group is provided by a noncommutative algebra, the noncommutative geometry presented by Connes [Con] plays a basic role like the differential geometry in the usual gauge theory. Following the general ideas of Connes, Woronowicz [Wor1] [Wor2] developed the framework of the noncommutative differential calculus. He introduced the bimodule over the quantum group, and presented various theorems concerning the differential forms and exterior derivative. Manin [Manin] suggested a general construction of quantum groups as linear transformations upon the quantum superplane. The differential calculus on the quantum hyperplane was developed by Wess and Zumino [WZ]. There have been a lot of papers treating the differential calculus on quantum groups and the deformed gauge theories from various viewpoints [Ber] [Jur] [Zum2] [CSWW] [IP] [AC] [Cas] [SW] [FP] [Isa]. Here we would like to emphasize three papers on the noncommutative differential geometry and deformation of BRST algebra.

Aschieri and Castellani [AC] gave a pedagogical introduction to the differential calculus on quantum groups by stressing at all stages the connection with the classical case ($q \rightarrow 1$). As an example, they gave the explicit forms of some matrices appearing in the bicovariant differential calculus on $SU_q(2)$. CSWW [CSWW] presented a systematic construction of bicovariant bimodules on the quantum groups $SU_q(N)$ and $SO_q(N)$ by using the $\hat{R}_q$ matrix. They described the conjugate of the fundamental representation for $SU_q(N)$ as antisymmetrized product of $(N - 1)$ fundamental representations, and showed the expressions appearing in the bicovariant differential calculus on $SU_q(N)$ both by formulas and by diagrams. On the other hand, the antisymmetrized product makes the calculation of explicit forms very complicated. In the later paper [Wat] Watamura investigated the q -deformation of BRST algebra for the quantum group $SU_q(2)$. Its generalization to $SU_q(N)$ depends on the explicit forms of the important quantities in the bicovariant differential calculus on $SU_q(N)$, such as the q -deformed structure constant $\mathbf{C}_{IJ}^K$, the q -deformed transposition operator Λ , and projection operator $\mathcal{P}_{Adj}$.

Based on the q -deformed BRST algebra presented by Watamura, we defined the q -deformed Killing form from the q -gauge covariant condition, and construct the second q -deformed Chern class, q -deformed Chern-Simons, and the cocycle hierarchy for $SU_q(2)$ in our previous paper [HHM]. In order to investigate the $SU_q(N)$ gauge theory we have to compute the explicit forms of the quantities in the bicovariant differential calculus on $SU_q(N)$.

In fact, the key for solving this problem is to change the description for the conjugate representation. As everyone knows, the conjugate of the fundamental representation in $SU(N)$ is equivalent to the antisymmetrized product of $(N - 1)$ fundamental representations, as used by [CSWW]. However, it is also equivalent to a basic highest weight representation described by the last fundamental dominant weight λ_{N-1} . The monoid $\epsilon^{a\bar{b}}$, that is an $N \times N$ matrix, plays a very important role in the explicit calculations. The monoid $\epsilon^{a\bar{b}}$, that seems to have some relation with the q -deformed Weyl element [KR], is proportional to the q -deformed Clebsch-Gordan matrix reducing the direct representation space of $\lambda_1 \otimes \lambda_{N-1}$ into the identity representation space [Ma], and provides the relations among the relevant $\hat{R}_q$ matrices. On the other hand, it also serves as the similarity transformation from the conjugate of the fundamental representation to the highest weight representation λ_{N-1} . From those $\hat{R}_q$ matrices and monoid we compute the explicit forms of the quantities appearing in the bicovariant differential calculus on $SU_q(N)$ so that we are able to generalize the q -deformed $SU_q(2)$ gauge theory to the quantum groups $SU_q(N)$. In other words, for the quantum groups $SU_q(N)$, we generalize the q -deformed BRST algebra, define the q -deformed Killing form, and construct the m -th q -deformed Chern class and q -deformed cocycle hierarchy. In [CSWW] and [CW] a similar matrix $\epsilon^{i[j]}$ was introduced, where $[j]$ is the short form for $(N - 1)$ antisymmetrized indices. However, our $\epsilon^{a\bar{b}}$ makes the calculations much simpler.

Recently, Isaev [Isa] discussed the q -deformed Chern characters where the base manifold is the q -deformed coset space $(GL_q(N + 1)/(GL_q(N) \otimes GL(1)))$. In the present paper, just like in [CSWW] and [AC], the spacetime is taken to be the ordinary commutative Minkowski spacetime, while the q -structure resides on the fiber, the gauge

potentials being non-commutating. In the theory there are two nilpotent operators: the BRST transformation δ and the derivative d , such that we can discuss the double cohomology and cocycle hierarchy. In terms of a q -deformed homotopy operator we are able to compute the q -deformed Chern-Simons Q_{2m-1} by the condition $dQ_{2m-1} = P_m$. As example, we write the explicit forms of the q -deformed Chern-Simons Q_3 and Q_5 explicitly.

The plan of this paper is as follows. In Sec. 2 we calculate the $\hat{R}_q$ matrices in the product representation spaces of $\lambda_1 \otimes \lambda_1$, $\lambda_{N-1} \otimes \lambda_{N-1}$, $\lambda_1 \otimes \lambda_{N-1}$, and $\lambda_{N-1} \otimes \lambda_1$, and discuss their main properties. The monoid $\epsilon^{a\bar{b}}$ is introduced to relate those $\hat{R}_q$ matrices. The algebra of functions on the quantum group $SU_q(N)$ is sketched in Sec. 3. It is proved that the monoid transforms the conjugate of the fundamental representation λ_1 into the highest weight representation λ_{N-1} . The generalized q -Pauli matrices are defined to separate the singlet from the adjoint components. In Sec. 4 we review the bicovariant differential calculus on $SU_q(N)$, and compute the explicit forms of some important quantities. Generalizing Watamura's investigation, the q -deformed BRST algebra for the quantum group $SU_q(N)$ is constructed in Sec. 5. From the condition $\delta P_m = 0$ and $dP_m = 0$, we define the q -deformed generalized Killing form and the m -th q -deformed Chern class P_m in Sec. 6. In Sec. 7, the q -deformed homotopy operator is introduced in $SU_q(N)$ to compute the q -deformed Chern-Simons. Furthermore, the q -deformed cocycle hierarchy and q -deformed Yang-Mills equation for $SU_q(N)$ gauge theory are obtained in Sec. 7 and Sec. 8. Just like in the case of $SU_q(2)$, the components of the identity and the adjoint representations are also separated in the q -deformed $SU_q(N)$ gauge theory, although they are mixed in the commutative relations of BRST algebra.

2. $\hat{R}_q$ Matrices and Monoid

In the quantum enveloping algebra $U_q A_{N-1}$ there are $(N-1)$ simple roots $\mathbf{r}_n$ and

$(N - 1)$ fundamental dominant weight λ_m , that are related by the Cartan matrix a_{mn} :

$$\begin{aligned} \mathbf{r}_n &= \lambda_m a_{mn}, \quad n, m = 1, 2, \dots, (N - 1) \\ a_{mn} &= d_m^{-1} (\mathbf{r}_m \cdot \mathbf{r}_n), \quad d_m = (\mathbf{r}_m \cdot \mathbf{r}_m) / 2 \end{aligned} \quad (2.1)$$

In the present paper, if without special notification, summation of the repeated indices is understood. To make formulas more symmetrical, we define:

$$\lambda_0 = \lambda_N = 0 \quad (2.2)$$

A highest weight representation is denoted by its highest weight, that is a positive integral combination of λ_j . The states in the representation are described by their weights that are the integral combinations of λ_j . In this paper we are interested in only eight representations: the basic representations $\lambda_1, \lambda_2, \lambda_{N-2}$ and λ_{N-1} , the symmetrical tensor representations $2\lambda_1$ and $2\lambda_{N-1}$, the adjoint representation $\lambda_1 + \lambda_{N-1}$, and the identity representation with the highest weight $\mathbf{0}$. Sometimes, the representations λ_2 and λ_{N-2} are also called the antisymmetrical tensor representations.

The fundamental representation λ_1 is N -dimensional. The states in this representation are described by their weight $(\lambda_a - \lambda_{a-1})$. For simplicity, we enumerate the states by one index a as usual:

$$a \rightarrow \lambda_a - \lambda_{a-1}, \quad a = 1, 2, \dots, N \quad (2.3)$$

The conjugate of the representation λ_1 is equivalent to the representation λ_{N-1} , where the states have the weights $\lambda_{a-1} - \lambda_a$ and are enumerated by one index $\bar{a}$:

$$\bar{a} \equiv -a \rightarrow \lambda_{a-1} - \lambda_a, \quad \bar{a} = \bar{N}, \dots, \bar{2}, \bar{1}, \quad (2.4)$$

In the present paper, if without special notification, the small Latin letter except for n and m , such as a and i , runs over $1, 2, \dots, N$.

The standard method for calculating the solutions $\hat{R}_q$ of the simple Yang-Baxter equation [Res] [Ma] is to expand it by the projection operators that are the products of two quantum Clebsch-Gordan matrices. The calculation method for $\hat{R}_q$ was described in the textbook [Ma] in detail. In order to fit the usual notation in the theory of the

quantum groups, the solution $\hat{R}_q$ here is related to the solution $\check{R}_q$ used in the book [Ma] as follows:

$$\hat{R}_q = q \check{R}_q^{-1} \quad (2.5)$$

In the direct product space of $\lambda_1 \otimes \lambda_1$ the solution of the simple Yang-Baxter equation is:

$$\left(\hat{R}_q^{\lambda_1 \lambda_1}\right)_{cd}^{ab} = q \left(\mathcal{P}_{2\lambda_1}^{\lambda_1 \lambda_1}\right)_{cd}^{ab} - q^{-1} \left(\mathcal{P}_{\lambda_2}^{\lambda_1 \lambda_1}\right)_{cd}^{ab} \quad (2.6)$$

Similarly, in the direct product spaces of $\lambda_{N-1} \otimes \lambda_{N-1}$, $\lambda_1 \otimes \lambda_{N-1}$, and $\lambda_{N-1} \otimes \lambda_1$, the solutions of the simple Yang-Baxter equation are as follows, respectively:

$$\begin{aligned} \left(\hat{R}_q^{\lambda_{N-1} \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} &= q \left(\mathcal{P}_{2\lambda_{N-1}}^{\lambda_{N-1} \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} - q^{-1} \left(\mathcal{P}_{\lambda_{N-2}}^{\lambda_{N-1} \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} \\ \left(\hat{R}_q^{\lambda_1 \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} &= q \left(\mathcal{P}_{\lambda_1 + \lambda_{N-1}}^{\lambda_1 \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} + (-q)^{1-N} \left(\mathcal{P}_0^{\lambda_1 \lambda_{N-1}}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} \\ \left(\hat{R}_q^{\lambda_{N-1} \lambda_1}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} &= q \left(\mathcal{P}_{\lambda_1 + \lambda_{N-1}}^{\lambda_{N-1} \lambda_1}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} + (-q)^{1-N} \left(\mathcal{P}_0^{\lambda_{N-1} \lambda_1}\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} \end{aligned} \quad (2.7)$$

The superscripts, for example $\lambda_1 \lambda_1$, have been implied in the super- and sub-scripts ab and cd , and can be neglected. Now, through straightforward calculation, we obtain:

$$\begin{aligned} \left(\hat{R}_q\right)_{cd}^{ab} &= \left(\hat{R}_q\right)_{ab}^{cd} = \left(\hat{R}_q\right)_{\bar{d}\bar{c}}^{\bar{b}\bar{a}} = \left(\hat{R}_q\right)_{\bar{b}\bar{a}}^{\bar{d}\bar{c}} \\ &= \begin{cases} q & \text{when } a = b = c = d \\ \lambda & \text{when } a = c < b = d \\ 1 & \text{when } a = d \neq b = c \\ 0 & \text{the else cases} \end{cases} \\ \left(\hat{R}_q^{-1}\right)_{cd}^{ab} &= \left(\hat{R}_q^{-1}\right)_{ab}^{cd} = \left(\hat{R}_q^{-1}\right)_{\bar{d}\bar{c}}^{\bar{b}\bar{a}} = \left(\hat{R}_q^{-1}\right)_{\bar{b}\bar{a}}^{\bar{d}\bar{c}} \\ &= \begin{cases} q^{-1} & \text{when } a = b = c = d \\ -\lambda & \text{when } a = c > b = d \\ 1 & \text{when } a = d \neq b = c \\ 0 & \text{the else cases} \end{cases} \\ \left(\hat{R}_q\right)_{\bar{c}\bar{d}}^{\bar{a}\bar{b}} &= \left(\hat{R}_q\right)_{\bar{a}\bar{b}}^{c\bar{d}} = \begin{cases} 1 & \text{when } a = b = c = d \\ -q\lambda(-q)^{c-b} & \text{when } a = b > c = d \\ q & \text{when } a = d \neq b = c \\ 0 & \text{the else cases} \end{cases} \end{aligned} \quad (2.8)$$

$$\left(\hat{R}_q^{-1}\right)_{c\bar{d}}^{\bar{a}b} = \left(\hat{R}_q^{-1}\right)_{\bar{a}b}^{c\bar{d}} = \begin{cases} 1 & \text{when } a = b = c = d \\ q^{-1}\lambda(-q)^{c-b} & \text{when } a = b < c = d \\ q^{-1} & \text{when } a = d \neq b = c \\ 0 & \text{the else cases} \end{cases}$$

Hereafter, λ and the q -number $[m]$ are defined as follows:

$$\begin{aligned} \lambda &= q - q^{-1}, & \lambda^2 + 4 &= [2]^2, & \lambda^2 + 3 &= [3] \\ \lambda^2 + 2 &= \frac{[4]}{[2]}, & \lambda^2 + 1 &= \frac{[6]}{[3][2]} \\ [m] &= \frac{q^m - q^{-m}}{q - q^{-1}} \end{aligned} \quad (2.9)$$

Remind that in the direct product space of $\lambda_1 \otimes \lambda_{N-1}$ there is a subspace belonging to the identity representation. We define the monoid $\epsilon^{a\bar{b}}$, that is proportional to the quantum Clebsch-Gordan coefficients ([Ma] P.157) reducing the product space into the subspace, as follows:

$$\begin{aligned} \epsilon^{a\bar{b}} &= (-1)^{N-1} \epsilon_{a\bar{b}} = \delta_{ab} (-1)^{N-a} q^{a-(N+1)/2} \\ \epsilon_{\bar{b}a} &= (-1)^{N-1} \epsilon^{\bar{b}a} = \delta_{ab} (-1)^{N-a} q^{-a+(N+1)/2} \\ \epsilon^{a\bar{b}} &= (-1)^{N-1} D^a_c \epsilon^{\bar{b}c} = \epsilon_{\bar{b}c} D^c_a \end{aligned} \quad (2.10)$$

$$\begin{aligned} \epsilon^{a\bar{b}} \epsilon_{\bar{b}c} &= \delta_c^a, & \epsilon_{\bar{a}b} \epsilon^{b\bar{c}} &= \delta_a^c \\ \epsilon^{a\bar{b}} \epsilon_{c\bar{b}} &= (-1)^{N-1} D^a_c, & \epsilon_{\bar{b}a} \epsilon^{\bar{b}c} &= (-1)^{N-1} (D^{-1})^c_a \end{aligned} \quad (2.11)$$

where the diagonal matrix D is related to double antipode action (see Sec. 3):

$$D^a_b = \delta_b^a q^{-N+2a-1} = (-1)^{N-1} \epsilon^{a\bar{d}} \epsilon_{b\bar{d}} \quad (2.12)$$

Four $\hat{R}_q$ matrices can be related by $\epsilon^{a\bar{b}}$ matrices:

$$\begin{aligned} \left(\hat{R}_q^{\pm 1}\right)_{\bar{a}b}^{c\bar{d}} &= q^{\pm 1} \epsilon_{\bar{a}r} \left(\hat{R}_q^{\mp 1}\right)_{bs}^{rc} \epsilon^{s\bar{d}} = q^{\pm 1} \epsilon^{c\bar{r}} \left(\hat{R}_q^{\mp 1}\right)_{\bar{r}\bar{a}}^{\bar{d}\bar{s}} \epsilon_{\bar{s}b} \\ \left(\hat{R}_q^{\pm 1}\right)_{c\bar{d}}^{\bar{a}b} &= q^{\pm 1} \epsilon^{\bar{a}r} \left(\hat{R}_q^{\mp 1}\right)_{rc}^{bs} \epsilon_{s\bar{d}} = q^{\pm 1} \epsilon_{c\bar{r}} \left(\hat{R}_q^{\mp 1}\right)_{\bar{d}\bar{s}}^{\bar{r}\bar{a}} \epsilon^{\bar{s}b} \end{aligned} \quad (2.13)$$

$\hat{R}_q$ matrix satisfies some important relations [Res]:

$$\left(\hat{R}_q^{\pm 1}\right)_{bd}^{ac} D^d_c = q^{\pm N} \delta_b^a \quad (2.14)$$

$$D^a_r D^c_s \left(\hat{R}_q^{\pm 1}\right)_{bd}^{rs} = \left(\hat{R}_q^{\pm 1}\right)_{rs}^{ac} D^r_b D^s_d \quad (2.15)$$

3. Algebra of Functions on the Quantum Group $SU_q(N)$

A quantum group is introduced as the non-commutative and non-cocommutative Hopf algebra $\mathcal{A} = Fun_q(G)$ obtained by continuous deformations of the Hopf algebra of the function of a Lie group. The associative algebra $\mathcal{A}$, the algebra of functions on the quantum group, is freely generated by non-commutating matrix entries T_b^a satisfying the relation:

$$\left(\hat{R}_q\right)_{rs}^{ab} T_c^r T_d^s = T_r^a T_s^b \left(\hat{R}_q\right)_{cd}^{rs} \quad (3.1)$$

where $\left(\hat{R}_q\right)_{cd}^{ab}$ is given in (2.6).

T_b^a , the elements of the fundamental representation of quantum group $SU_q(N)$, satisfy the Hopf algebraic relations:

$$\begin{aligned} \Delta(T_b^a) &= T_d^a \otimes T_b^d, \quad \epsilon(T_b^a) = \delta_b^a \\ \kappa(T_r^a) T_b^r &= T_r^a \kappa(T_b^r) = \delta_b^a, \quad \kappa^2(T_b^a) = D_c^a T_d^c (D^{-1})_b^d \end{aligned} \quad (3.2)$$

where the diagonal matrix D is given in (2.12).

The q -determinant $det_q T$ is commutant with any element T_b^a . For quantum group $SU_q(N)$ we have:

$$\begin{aligned} det_q T &= \sum_P (-q)^{\ell(P)} T_{p_1}^1 \cdots T_{p_N}^N = 1, \\ q^* &= q, \quad (T_b^a)^* = \kappa(T_a^b) \end{aligned} \quad (3.3)$$

where $\ell(P)$ is the minimal number of inversions in the permutation P . $\kappa(T_b^a)$ can be expressed as antisymmetrized product of $(N-1)$ T_b^a [CSWW]. On the other hand, the conjugate of the fundamental representation λ_1 is equivalent to the representation λ_{N-1} . Define:

$$T_{\bar{b}}^{\bar{a}} = \epsilon_{\bar{b}r} \kappa(T_s^r) \epsilon^{s\bar{a}}, \quad \kappa(T_a^b) = \epsilon^{b\bar{r}} T_{\bar{r}}^{\bar{s}} \epsilon_{\bar{s}a} \quad (3.4)$$

For $SU_q(2)$, $\lambda_1 = \lambda_{N-1}$. In this case the fundamental representation becomes self-conjugate:

$$T_{\bar{b}}^{\bar{a}} = T_{3-b}^{3-a}$$

From (3.1), (3.2) and (2.13) we can show that $T_{\bar{b}}^{\bar{a}} \in \mathcal{A}$ defined in (3.4) belongs to the representation λ_{N-1} .

Proposition 1. $T_{\bar{b}}^{\bar{a}} \in \mathcal{A}$ defined in (3.4) satisfy:

$$\begin{aligned} \left(\hat{R}_q \right)_{r\bar{s}}^{\bar{a}b} T_c^r T_{\bar{d}}^{\bar{s}} &= T_{\bar{r}}^{\bar{a}} T_s^b \left(\hat{R}_q \right)_{cd}^{\bar{r}s} \\ \left(\hat{R}_q \right)_{\bar{r}s}^{\bar{a}b} T_{\bar{c}}^{\bar{r}} T_d^s &= T_r^a T_{\bar{s}}^{\bar{b}} \left(\hat{R}_q \right)_{\bar{c}d}^{r\bar{s}} \\ \left(\hat{R}_q \right)_{\bar{r}\bar{s}}^{\bar{a}\bar{b}} T_{\bar{c}}^{\bar{r}} T_{\bar{d}}^{\bar{s}} &= T_{\bar{r}}^{\bar{a}} T_{\bar{s}}^{\bar{b}} \left(\hat{R}_q \right)_{\bar{c}\bar{d}}^{\bar{r}\bar{s}} \end{aligned} \quad (3.5)$$

Proof. From (3.1) and (3.2) we have

$$\kappa(T_s^a) \left(\hat{R}_q \right)_{rd}^{sb} T_c^r = T_s^b \left(\hat{R}_q \right)_{cr}^{as} \kappa(T_d^r)$$

In terms of (2.13) and (3.4) we obtain:

$$\begin{aligned} \kappa(T_s^a) \left(\hat{R}_q \right)_{rd}^{sb} T_c^r &= \epsilon^{a\bar{t}} T_{\bar{t}}^{\bar{u}} \epsilon_{\bar{u}s} \left(\hat{R}_q \right)_{rd}^{sb} T_c^r \\ &= q \epsilon^{a\bar{t}} \left\{ T_{\bar{t}}^{\bar{u}} \left(\hat{R}_q^{-1} \right)_{\bar{u}r}^{b\bar{v}} T_c^r \right\} \epsilon_{\bar{v}d} \\ T_s^b \left(\hat{R}_q \right)_{cr}^{as} \kappa(T_d^r) &= T_s^b \left(\hat{R}_q \right)_{cr}^{as} \epsilon^{r\bar{u}} T_{\bar{u}}^{\bar{v}} \epsilon_{\bar{v}d} \\ &= q \epsilon^{a\bar{t}} \left\{ T_s^b \left(\hat{R}_q^{-1} \right)_{\bar{t}c}^{s\bar{u}} T_{\bar{u}}^{\bar{v}} \right\} \epsilon_{\bar{v}d} \end{aligned}$$

Comparing the quantities in the brackets, we get the first relation in (3.5). The proof of the other two relations in (3.5) can be performed analogously. Q.E.D.

The direct product of T_c^a and $T_{\bar{d}}^{\bar{b}}$ spans the mixed space of the adjoint and identity representations:

$$T_c^a T_{\bar{d}}^{\bar{b}} = T_c^a \epsilon_{\bar{d}r} \kappa(T_s^r) \epsilon^{s\bar{b}} \quad (3.6)$$

In order to separate the singlet and the adjoint components in $T_c^a T_{\bar{d}}^{\bar{b}}$, we define the generalized q -Pauli matrices, that obviously have to be proportional to the quantum Clebsch-Gordan coefficients:

$$\begin{aligned} (\sigma^0)_{a\bar{b}} &= q^2 (\sigma_0)^{a\bar{b}} = -q [N]^{-1/2} \epsilon^{a\bar{b}} \\ (\sigma^{i\bar{j}})_{a\bar{b}} &= (\sigma_{i\bar{j}})^{a\bar{b}} = (-1)^{N-b} \delta_{ai} \delta_{b\bar{j}}, \quad i \neq j \\ (\sigma^{j\bar{j}})_{a\bar{b}} &= \delta_{ab} \begin{cases} (-1)^{N+a} q^{a-j+(N+1)/2} \{[j-1][j]\}^{-1/2} & \text{when } a < j \\ (-1)^{N+j+1} q^{-j+(N+1)/2} \{[j-1]/[j]\}^{1/2} & \text{when } a = j, \quad j \geq 2 \\ 0 & \text{when } a > j \end{cases} \\ (\sigma_{j\bar{j}})^{a\bar{b}} &= \delta_{ab} \begin{cases} (-1)^{N+a} q^{a+j-(N+3)/2} \{[j-1][j]\}^{-1/2} & \text{when } a < j \\ (-1)^{N+j+1} q^{j-(N+3)/2} \{[j-1]/[j]\}^{1/2} & \text{when } a = j, \quad j \geq 2 \\ 0 & \text{when } a > j \end{cases} \end{aligned} \quad (3.7)$$

where the repeated j is not summed. In the present paper, we describe the adjoint components by a capital Latin, such as I , that runs over $(i\bar{j})$ where $i \neq j$, and $(j\bar{j})$ where $j \geq 2$, and describe both singlet and adjoint components by a capital Latin with a hat, like $\hat{I}$, that runs over 0 and I . In this notation, we have:

$$(\sigma^{\hat{I}})_{a\bar{b}} (\sigma_j)^{a\bar{b}} = \delta_{\hat{I}j}, \quad (\sigma^{\hat{I}})_{a\bar{b}} (\sigma_{\hat{I}})^{c\bar{d}} = \delta_a^c \delta_b^d \quad (3.8)$$

Sometime, another set of q -Pauli matrices are useful:

$$\begin{aligned} (\sigma^{\hat{I}})_a^b &= (\sigma^{\hat{I}})_{a\bar{d}} \epsilon^{b\bar{d}}, & (\sigma_{\hat{I}})_b^a &= (\sigma_{\hat{I}})^{a\bar{d}} \epsilon_{\bar{d}b} \\ (\sigma^{\hat{I}})_a^b (\sigma_j)_b^a &= \delta_{\hat{I}j}, & (\sigma^{\hat{I}})_a^b (\sigma_{\hat{I}})_d^c &= \delta_a^c \delta_b^d \end{aligned} \quad (3.9)$$

$$\begin{aligned} (\sigma^0)_a^b &= -q [N]^{-1/2} D^b_a, & (\sigma_0)_b^a &= -q^{-1} [N]^{-1/2} \delta_b^a \\ (\sigma^{i\bar{j}})_a^b &= \delta_a^i \delta_b^j q^{b-(N+1)/2}, & (\sigma_{i\bar{j}})_b^a &= \delta_i^a \delta_j^b q^{-b+(N+1)/2}, \quad i \neq j \\ (\sigma^{j\bar{j}})_a^b &= \delta_{ab} \begin{cases} q^{2a-j} \{[j-1][j]\}^{-1/2} & \text{when } a < j \\ -\{[j-1]/[j]\}^{1/2} & \text{when } a = j, \quad j \geq 2 \\ 0 & \text{when } a > j \end{cases} \\ (\sigma_{j\bar{j}})_b^a &= \delta_{ab} \begin{cases} q^{j-1} \{[j-1][j]\}^{-1/2} & \text{when } a < j \\ -q^{-1} \{[j-1]/[j]\}^{1/2} & \text{when } a = j, \quad j \geq 2 \\ 0 & \text{when } a > j \end{cases} \end{aligned} \quad (3.10)$$

where the repeated j is not summed. When $q \rightarrow 1$, both $(\sigma^{\hat{I}})_a^b$ and $(\sigma_{\hat{I}})_d^c$ tend to the usual generalized Pauli matrices. Now, it can be seen that the singlet and the adjoint components in $T_c^a T_{\bar{d}}^{\bar{b}}$ are separated explicitly:

$$\begin{aligned} M_{\hat{j}}^{\hat{I}} &= (\sigma^{\hat{I}})_{a\bar{b}} T_c^a T_{\bar{d}}^{\bar{b}} (\sigma_j)^{c\bar{d}} = (\sigma^{\hat{I}})_a^d T_c^a \kappa(T_d^b) (\sigma_j)_b^c \\ M_0^0 &= \mathbf{1}, \quad M_0^I = M_I^0 = 0 \\ \kappa^2(M_{\hat{j}}^{\hat{I}}) &= D_{\hat{K}}^{\hat{I}} M_{\hat{L}}^{\hat{K}} (D^{-1})_{\hat{J}}^{\hat{L}} \\ D_0^0 &= 1, \quad D_{(i\bar{j})}^{(i\bar{j})} = q^{2(i-j)} \end{aligned} \quad (3.11)$$

where D is a diagonal matrix.

The linear functionals $(L^{\pm})_b^a$, defined by their values on the entries T_b^a , belong to the dual Hopf algebra $\mathcal{A}'$ [FRT]:

$$(L^+)_b^a (T_d^c) = q^{-1/N} (\hat{R}_q)_{db}^{ac}, \quad (L^-)_b^a (T_d^c) = q^{1/N} (\hat{R}_q^{-1})_{db}^{ac} \quad (3.12)$$

$$\begin{aligned}\Delta'((L^\pm)^a_b) &= (L^\pm)^a_c \otimes (L^\pm)^c_b, \quad \epsilon'((L^\pm)^a_b) = \delta_b^a \\ \kappa'((L^\pm)^a_c) (L^\pm)^c_b &= \delta_b^a = (L^\pm)^a_c \kappa'((L^\pm)^c_b)\end{aligned}\tag{3.13}$$

$$\begin{aligned}(\hat{R}_q)^{ab}_{rs} (L^\pm)^s_d (L^\pm)^r_c &= (L^\pm)^b_s (L^\pm)^a_r (\hat{R}_q)^{rs}_{cd} \\ (\hat{R}_q)^{ab}_{rs} (L^+)^s_d (L^-)^r_c &= (L^+)^b_s (L^-)^a_r (\hat{R}_q)^{rs}_{cd}\end{aligned}\tag{3.14}$$

From (3.12) we have:

$$\begin{aligned}\kappa'((L^\pm)^a_b) (T^c_d) &= q^{\pm 1/N} (\hat{R}_q^{\mp 1})^{ca}_{bd} \\ ((L^\pm)^a_b) (T^{\bar{c}}_{\bar{d}}) &= q^{\pm 1/N} \epsilon_{\bar{d}r} (\hat{R}_q^{\mp 1})^{ra}_{bs} \epsilon^{s\bar{c}} = q^{\mp(1-1/N)} (\hat{R}_q^{\pm 1})^{a\bar{c}}_{\bar{d}b} \\ \kappa'((L^\pm)^a_b) (T^{\bar{c}}_{\bar{d}}) &= q^{\mp 1/N} \epsilon_{\bar{d}r} D^r_u (\hat{R}_q^{\pm 1})^{au}_{vb} (D^{-1})^v_s \epsilon^{s\bar{c}}\end{aligned}\tag{3.15}$$

4. Bicovariant Differential Calculus on Quantum Groups $SU_q(N)$

[CSWW] constructed the bimodule Γ for the quantum groups $SU_q(N)$ explicitly. In order to avoid confusion with the spacetime derivative, following Watamura's notation [Wat] in the theory of q -deformed BRST algebra, we call the first-order differential operator on $\mathcal{A}$ the BRST transformation operator, denoted by δ :

$$\begin{aligned}\delta : \mathcal{A} &\rightarrow \Gamma \\ \rho &= \sum \alpha \delta \beta \in \Gamma, \quad \text{if } \rho \in \Gamma\end{aligned}\tag{4.1}$$

where $\alpha, \beta \in \mathcal{A}$.

A left action Δ_L and a right action Δ_R of the quantum group on Γ are defined as follows:

$$\begin{aligned}\Delta_L : \Gamma &\rightarrow \mathcal{A} \otimes \Gamma, \quad \Delta_L(\alpha \delta \beta) = \Delta(\alpha) (id \otimes \delta) \Delta(\beta) \\ \Delta_R : \Gamma &\rightarrow \Gamma \otimes \mathcal{A}, \quad \Delta_R(\alpha \delta \beta) = \Delta(\alpha) (\delta \otimes id) \Delta(\beta)\end{aligned}\tag{4.2}$$

The tensor product between elements $\rho, \rho' \in \Gamma$ is defined to have the properties:

$$\begin{aligned}(\rho \alpha) \otimes \rho' &= \rho \otimes (\alpha \rho') \\ \alpha(\rho \otimes \rho') &= (\alpha \rho) \otimes \rho', \quad (\rho \otimes \rho')\alpha = \rho \otimes (\rho' \alpha)\end{aligned}\tag{4.3}$$

The left and right actions on $\Gamma \otimes \Gamma$ are defined by:

$$\begin{aligned}\Delta_L : \Gamma \otimes \Gamma &\rightarrow \mathcal{A} \otimes \Gamma \otimes \Gamma \\ \Delta_R : \Gamma \otimes \Gamma &\rightarrow \Gamma \otimes \Gamma \otimes \mathcal{A}\end{aligned}\tag{4.4}$$

For example,

$$\begin{aligned}\Delta_L(\rho_1) &= \alpha_1 \otimes \rho'_1, & \Delta_L(\rho_2) &= \alpha_2 \otimes \rho'_2 \\ \Delta_L(\rho_1 \otimes \rho_2) &= \alpha_1 \alpha_2 \otimes \rho'_1 \otimes \rho'_2\end{aligned}\tag{4.5}$$

From the definition we have:

$$\begin{aligned}(\epsilon \otimes id) \Delta_L(\rho) &= \rho, & (id \otimes \epsilon) \Delta_R(\rho) &= \rho \\ (\Delta \otimes id) \Delta_L &= (id \otimes \Delta_L) \Delta_L, & (id \otimes \Delta) \Delta_R &= (\Delta_R \otimes id) \Delta_R \\ (id \otimes \Delta_R) \Delta_L &= (\Delta_L \otimes id) \Delta_R\end{aligned}\tag{4.6}$$

[CSWW] constructed the fundamental bicovariant bimodule of $SU_q(N)$:

$$\begin{aligned}\Delta_R(\eta^a) &= \eta^a \otimes \mathbf{1}, & \Delta_L(\eta^a) &= T^a_b \otimes \eta^b \\ \eta^a \alpha &= (\alpha * f^a_b) \eta^b, & \alpha \eta^a &= \eta^b (\alpha * f^a_b \circ \kappa)\end{aligned}\tag{4.7}$$

where $(\alpha * f^a_b) \equiv (f^a_b \otimes id) \Delta \alpha$.

Applying the $*$ -operation on (4.7), [CSWW] obtained

$$\begin{aligned}(\eta^a)^* &\equiv \bar{\eta}_a, & \Delta_R(\bar{\eta}_a) &= \bar{\eta}_a \otimes \mathbf{1}, & \Delta_L(\bar{\eta}_a) &= \kappa(T^b_a) \otimes \bar{\eta}_b \\ \bar{\eta}_a \alpha &= (\alpha * \bar{f}^b_a) \bar{\eta}_b, & \alpha \bar{\eta}_a &= \bar{\eta}_b (\alpha * \bar{f}^b_a \circ \kappa)\end{aligned}\tag{4.8}$$

From the consistency with the commutation relation of the generators and the requirement for $SU_q(N)$ that the q -determinant of T^a_b commutes with any element in Γ , [CSWW] found that there are two independent functionals such that the following constructions are performed in a completely parallel way. Following [CSWW] we choose one of them as follows:

$$f^a_b = (L^+)^a_b, \quad \bar{f}^a_b = \kappa'^{-1} \left((L^-)^a_b \right)\tag{4.9}$$

Transforming the bases $\bar{\eta}_a$ by the monoid $\epsilon^{a\bar{b}}$, we obtain the bases $\eta^{\hat{I}}$ for the mixed representation of identity and adjoint representations:

$$\begin{aligned}\bar{\eta}^{\bar{b}} &= \bar{\eta}_a \epsilon^{a\bar{b}}, & \bar{\eta}_a &= \bar{\eta}^{\bar{b}} \epsilon_{\bar{b}a} \\ \eta^{\hat{I}} &= (\sigma^{\hat{I}})_{a\bar{b}} \eta^a \bar{\eta}^{\bar{b}} = (\sigma^{\hat{I}})_a^b \eta^a \bar{\eta}_b\end{aligned}\tag{4.10}$$

In the commutative relations of the q -deformed BRST algebra the components of identity and adjoint representations are mixed. For the bases $\eta^{\hat{I}}$ we have

$$\begin{aligned}\Delta_R(\eta^{\hat{J}}) &= \eta^{\hat{J}} \otimes \mathbf{1}, & \Delta_L(\eta^{\hat{J}}) &= M^{\hat{J}}_{\hat{K}} \otimes \eta^{\hat{K}} \\ \alpha \eta^{\hat{J}} &= \eta^{\hat{K}} (\alpha * L^{\hat{J}}_{\hat{K}}), & \alpha \in \mathcal{A}, & L^{\hat{J}}_{\hat{K}} \in \mathcal{A}'\end{aligned}\tag{4.11}$$

where

$$\begin{aligned}
L_{\hat{I}}^{\hat{J}} &= (\sigma^{\hat{J}})_a^d \kappa'((L^+)^a_b) (L^-)^c_d (\sigma_{\hat{I}})^b_c \\
&= (\sigma^{\hat{J}})_{a\bar{r}} \left\{ \epsilon^{d\bar{r}} \kappa'((L^+)^a_b) (L^-)^c_d \epsilon_{\bar{s}c} \right\} (\sigma_{\hat{I}})^{b\bar{s}} \\
L_{\hat{I}}^{\hat{J}}(ab) &= L_{\hat{I}}^{\hat{K}}(a) L_{\hat{K}}^{\hat{J}}(b), \quad L_{\hat{I}}^{\hat{J}}(\mathbf{1}) = \delta_{\hat{I}}^{\hat{J}} \\
(\rho * L_{\hat{I}}^{\hat{J}}) &= (L_{\hat{I}}^{\hat{J}} \otimes id) \Delta_L(\rho)
\end{aligned} \tag{4.12}$$

Note that:

$$\Delta_L(\eta^0) = \mathbf{1} \otimes \eta^0, \quad \Delta_R(\eta^0) = \eta^0 \otimes \mathbf{1} \tag{4.13}$$

The bases of the left-invariant element of Γ are easy to be calculated from $\eta^{\hat{J}}$:

$$\omega^{\hat{J}} = \kappa(M_{\hat{K}}^{\hat{J}}) \eta^{\hat{K}}, \quad \Delta_L(\omega^{\hat{J}}) = \mathbf{1} \otimes \omega^{\hat{J}}, \quad \Delta_R(\omega^{\hat{J}}) = \omega^{\hat{K}} \otimes \kappa(M_{\hat{K}}^{\hat{J}}) \tag{4.14}$$

As the analogue of the ordinary permutation operator, a bimodule automorphism Λ in $\Gamma \otimes \Gamma$ is defined by:

$$\begin{aligned}
\Lambda(\omega^{\hat{J}} \otimes \eta^{\hat{K}}) &= \eta^{\hat{K}} \otimes \omega^{\hat{J}} \\
\Lambda(\alpha \tau) &= \alpha \Lambda(\tau), \quad \Lambda(\tau \alpha) = \Lambda(\tau) \alpha, \quad \alpha \in \mathcal{A}, \quad \tau \in \Gamma \otimes \Gamma
\end{aligned} \tag{4.15}$$

Thus, we have

$$\Lambda(\eta^{\hat{I}} \otimes \eta^{\hat{J}}) = \Lambda^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} \eta^{\hat{K}} \otimes \eta^{\hat{L}}, \quad \Lambda^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} = L_{\hat{K}}^{\hat{J}}(M_{\hat{L}}^{\hat{I}}) \tag{4.16}$$

In terms of (2.13), (3.12) and (3.15) we are able to compute $\Lambda^{\hat{I}\hat{J}}_{\hat{K}\hat{L}}$ explicitly:

$$\begin{aligned}
(\sigma_{\hat{I}})^{a\bar{b}} (\sigma_{\hat{J}})^{c\bar{d}} \Lambda^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} (\sigma_{\hat{K}})_{i\bar{j}} (\sigma_{\hat{L}})_{k\bar{\ell}} &= (\sigma_{\hat{K}})_{i\bar{j}} L_{\hat{K}}^{\hat{S}}(T_k^a) L_{\hat{S}}^{\hat{J}}(T_{\bar{\ell}}^{\bar{b}}) (\sigma_{\hat{J}})^{c\bar{d}} \\
&= \kappa'((L^+)^r_i) (T_u^a) \epsilon_{\bar{j}t} (L^-)^t_w (T_k^u) \epsilon^{w\bar{s}} \kappa'((L^+)^c_r) (T_{\bar{v}}^{\bar{b}}) \epsilon_{\bar{s}x} (L^-)^x_y (T_{\bar{\ell}}^{\bar{v}}) \epsilon^{y\bar{d}} \\
&= q^{1/N} (\hat{R}_q^{-1})_{iu}^{ar} \epsilon_{\bar{j}t} q^{1/N} (\hat{R}_q^{-1})_{kw}^{tu} \epsilon^{w\bar{s}} q^{-1/N} \epsilon_{\bar{v}r'} D_{s'}^{r'} (\hat{R}_q)^{cs'}_{u'r} (D^{-1})^{u'}_{v'} \epsilon^{v'\bar{b}} \\
&\quad \cdot \epsilon_{\bar{s}x} q^{1-1/N} (\hat{R}_q^{-1})_{\bar{\ell}y}^{x\bar{v}} \epsilon^{y\bar{d}} \\
&= q \epsilon_{\bar{j}t} \epsilon_{s'\bar{v}} \epsilon^{\bar{b}u'} \epsilon^{y\bar{d}} (\hat{R}_q^{-1})_{iu}^{ar} (\hat{R}_q^{-1})_{kw}^{tu} (\hat{R}_q)^{cs'}_{u'r} (\hat{R}_q^{-1})_{\bar{\ell}y}^{w\bar{v}} \\
&= (\hat{R}_q^{-1})_{iu}^{ar} (\hat{R}_q)^{u\bar{s}}_{\bar{j}k} (\hat{R}_q^{-1})_{r\bar{v}}^{\bar{b}c} (\hat{R}_q)^{\bar{v}\bar{d}}_{\bar{s}\bar{\ell}}
\end{aligned} \tag{4.17}$$

For given matrices $(\mathcal{P}_1)_{ij}^{ab}$ and $(\mathcal{P}_2)_{\bar{k}\bar{\ell}}^{\bar{c}\bar{d}}$ define a operator $(\mathcal{P}_1, \mathcal{P}_2)_{\hat{K}\hat{L}}^{\hat{I}\hat{J}}$ [CSWW]:

$$(\mathcal{P}_1, \mathcal{P}_2)_{\hat{K}\hat{L}}^{\hat{I}\hat{J}} = (\sigma^{\hat{I}})_{a\bar{b}} (\sigma^{\hat{J}})_{c\bar{d}} (\hat{R}_q^{-1})_{r\bar{v}}^{\bar{b}c} (X)_{iu}^{ar} (Y)_{\bar{s}\bar{\ell}}^{\bar{v}\bar{d}} (\hat{R}_q)^{u\bar{s}}_{\bar{j}k} (\sigma_{\hat{K}})^{i\bar{j}} (\sigma_{\hat{L}})^{k\bar{\ell}} \tag{4.18}$$

Hence, we obtain four projection operators, orthogonal to each other:

$$\begin{aligned}
& (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{2\lambda_{N-1}}), \quad (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{\lambda_{N-2}}), \quad (\mathcal{P}_{\lambda_2}, \mathcal{P}_{2\lambda_{N-1}}), \quad (\mathcal{P}_{\lambda_2}, \mathcal{P}_{\lambda_{N-2}}) \\
& \mathcal{P}_S \equiv (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{2\lambda_{N-1}}) + (\mathcal{P}_{\lambda_2}, \mathcal{P}_{\lambda_{N-2}}) \\
& \mathcal{P}_A \equiv (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{\lambda_{N-2}}) + (\mathcal{P}_{\lambda_2}, \mathcal{P}_{2\lambda_{N-1}}) \\
& \mathcal{P}_S + \mathcal{P}_A = \mathbf{1}
\end{aligned} \tag{4.19}$$

From (4.17), (2.6) and (2.7) we have

$$\begin{aligned}
\Lambda &= (\hat{R}^{-1}, \hat{R}) \\
&= (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{2\lambda_{N-1}}) - q^{-2} (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{\lambda_{N-2}}) - q^2 (\mathcal{P}_{\lambda_2}, \mathcal{P}_{2\lambda_{N-1}}) + (\mathcal{P}_{\lambda_2}, \mathcal{P}_{\lambda_{N-2}}) \\
\Lambda^{-1} &= (\hat{R}, \hat{R}^{-1}) \\
&= (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{2\lambda_{N-1}}) - q^2 (\mathcal{P}_{2\lambda_1}, \mathcal{P}_{\lambda_{N-2}}) - q^{-2} (\mathcal{P}_{\lambda_2}, \mathcal{P}_{2\lambda_{N-1}}) + (\mathcal{P}_{\lambda_2}, \mathcal{P}_{\lambda_{N-2}})
\end{aligned} \tag{4.20}$$

Therefore, $\Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}}$ satisfy the Yang-Baxter equation:

$$\Lambda_{\hat{L}\hat{S}}^{\hat{I}\hat{J}} \Lambda_{\hat{T}\hat{R}}^{\hat{S}\hat{K}} \Lambda_{\hat{P}\hat{Q}}^{\hat{L}\hat{T}} = \Lambda_{\hat{L}\hat{S}}^{\hat{J}\hat{K}} \Lambda_{\hat{P}\hat{T}}^{\hat{I}\hat{L}} \Lambda_{\hat{Q}\hat{R}}^{\hat{T}\hat{S}} \tag{4.21}$$

From (4.20) we know that the eigenvalues of Λ matrix are 1, $-q^2$ and $-q^{-2}$:

$$(\Lambda + q^2) (\Lambda + q^{-2}) (\Lambda - \mathbf{1}) = 0 \tag{4.22}$$

From the symmetry of the quantum Clebsch-Gordan coefficients ([Ma] P.156) we have:

$$(\Lambda^{-1})_{(a\bar{b})(c\bar{d})}^{(i\bar{j})(k\bar{\ell})} = \Lambda_{(d\bar{c})(b\bar{a})}^{(\ell\bar{k})(j\bar{i})} \tag{4.23}$$

Through direct calculation of (4.17), we obtain the non-vanishing components of $\Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}}$ as follows:

$$\begin{aligned}
\Lambda_{KL}^{IJ} &= (\Lambda^{-1})_{KL}^{IJ} = \delta_K^I \delta_L^J + f_P^{IJ} \bar{f}_{KL}^P, \quad \Lambda_{00}^{00} = (\Lambda^{-1})_{00}^{00} = 1 \\
\Lambda_{JK}^{I0} &= (\Lambda^{-1})_{JK}^{0I} = \lambda f_{JK}^I, \quad \Lambda_{0I}^{JK} = (\Lambda^{-1})_{0I}^{JK} = \lambda \bar{f}_I^{JK} \\
\Lambda_{J0}^{0I} &= (\Lambda^{-1})_{0J}^{I0} = \delta_J^I, \quad \Lambda_{0J}^{I0} = (\Lambda^{-1})_{0J}^{0I} = (\lambda^2 + 1) \delta_J^I
\end{aligned} \tag{4.24}$$

where f_{JK}^I and $\bar{f}_I^{JK}$ satisfy:

$$\begin{aligned}
f_{(i\bar{j})(k\bar{\ell})}^{(r\bar{s})} &= 0, \quad \bar{f}_{(r\bar{s})}^{(i\bar{j})(k\bar{\ell})} = 0, \\
&\text{if } \lambda_i - \lambda_{i-1} - \lambda_j + \lambda_{j-1} + \lambda_k - \lambda_{k-1} - \lambda_\ell + \lambda_{\ell-1} \neq \lambda_r - \lambda_{r-1} - \lambda_s + \lambda_{s-1} \\
f_{(i\bar{j})(k\bar{\ell})}^{(r\bar{s})} &= f_{(\ell\bar{k})(j\bar{i})}^{(s\bar{r})}, \quad \bar{f}_{(r\bar{s})}^{(i\bar{j})(k\bar{\ell})} = \bar{f}_{(s\bar{r})}^{(\ell\bar{k})(j\bar{i})}, \quad f_{RS}^I \bar{f}_J^{RS} = -(\lambda^2 + 2) \delta_J^I
\end{aligned} \tag{4.25}$$

The non-vanishing components of f_{JK}^I and $\bar{f}_I^{JK}$ are listed as follows. In the following (4.26) there is no summation for the repeated indices.

$$\begin{aligned} f_{(\bar{i}\bar{k})(k\bar{j})}^{(i\bar{j})} &= [N]^{-1/2} q^{-k-(N-3)/2}, & f_{(k\bar{j})(i\bar{k})}^{(i\bar{j})} &= - [N]^{-1/2} q^{k-(N-1)/2} \\ \bar{f}_{(i\bar{j})}^{(i\bar{k})(k\bar{j})} &= - [N]^{-1/2} q^{-k+(3N-1)/2}, & \bar{f}_{(i\bar{j})}^{(k\bar{j})(i\bar{k})} &= [N]^{-1/2} q^{k-(N+3)/2} \end{aligned} \quad (4.26a)$$

when $i \neq j \neq k \neq i$

$$\begin{aligned} f_{(k\bar{k})(i\bar{j})}^{(i\bar{j})} &= \begin{cases} -q^{-N} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = i < j \\ -q^{1-N} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} & \text{if } j < i = k \\ q^{2k-N-1} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} & \text{if } i < j = k \\ q^{2k-N} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = j < i \\ q^{k-N} ([N][k][k-1])^{-1/2} & \text{if } i < k < j \\ -q^{k-N} ([N][k][k-1])^{-1/2} & \text{if } j < k < i \end{cases} \quad (4.26b) \\ \bar{f}_{(i\bar{j})}^{(k\bar{k})(i\bar{j})} &= \begin{cases} q^{2N-2k} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = i < j \\ q^{2N-2k+1} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} - \lambda q^{N-k} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } j < i = k \\ -q^{-1} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} - \lambda q^{N-k} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } i < j = k \\ - \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = j < i \\ -q^{2N-k} ([N][k][k-1])^{-1/2} & \text{if } i < k < j \\ q^{-k} ([N][k][k-1])^{-1/2} & \text{if } j < k < i \\ -\lambda q^{N-k} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } i < j < k \\ & \text{or } j < i < k \end{cases} \quad (4.26c) \end{aligned}$$

$$f_{(i\bar{j})(j\bar{i})}^{(k\bar{k})} = \begin{cases} -q^{2-k-j} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = i < j \\ -q^{3-k-j} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} & \text{if } j < i = k \\ q^{i-k+1} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} & \text{if } i < j = k \\ q^{i-k+2} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = j < i \\ q^{i-j-k+2} ([N][k][k-1])^{-1/2} & \text{if } i < k < j \\ -q^{i-j-k+2} ([N][k][k-1])^{-1/2} & \text{if } j < k < i \end{cases} \quad (4.26d)$$

$$\bar{f}_{(k\bar{k})}^{(i\bar{j})(j\bar{i})} = \begin{cases} q^{N+k-j-2} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = i < j \\ q^{N+k-j-1} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} - \lambda q^{2k-j-2} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } j < i = k \\ -q^{i+k-N-3} \left(\frac{[k]}{[N][k-1]} \right)^{1/2} - \lambda q^{i-2} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } i < j = k \\ -q^{i+k-N-2} \left(\frac{[k-1]}{[N][k]} \right)^{1/2} & \text{if } k = j < i \\ -q^{i-j+k+N-2} ([N][k][k-1])^{-1/2} & \text{if } i < k < j \\ q^{i-j+k-N-2} ([N][k][k-1])^{-1/2} & \text{if } j < k < i \\ -\lambda q^{i-j+k-2} \left(\frac{[N]}{[k][k-1]} \right)^{1/2} & \text{if } i < j < k \\ & \text{or } j < i < k \end{cases} \quad (4.26e)$$

$$\begin{aligned} f_{(k\bar{k})(k\bar{k})}^{(k\bar{k})} &= \lambda q^{k-N} \left(\frac{[k][k-1]}{[N]} \right)^{1/2} \\ \bar{f}_{(k\bar{k})}^{(k\bar{k})(k\bar{k})} &= -\lambda q^{N-k} \left\{ \left(\frac{[N]}{[k][k-1]} \right)^{1/2} - [N-k] \left(\frac{[k-1]}{[N][k]} \right)^{1/2} \right\} \\ \bar{f}_{(k\bar{k})}^{(j\bar{j})(j\bar{j})} &= -\lambda q^{N-2j+k} \left(\frac{[N]}{[k][k-1]} \right)^{1/2}, \quad \text{if } j < k \\ \bar{f}_{(j\bar{j})}^{(j\bar{j})(k\bar{k})} &= \bar{f}_{(j\bar{j})}^{(k\bar{k})(j\bar{j})} = -\lambda q^{N-k} \left(\frac{[N]}{[k][k-1]} \right)^{1/2}, \quad \text{if } j < k \end{aligned} \quad (4.26f)$$

Substituting (4.24) into the Yang-Baxter equation (4.21) for two cases: $\hat{K} = \hat{P} = \hat{Q} = 0$,

and $\hat{J} = \hat{K} = \hat{P} = 0$, we obtain:

$$\begin{aligned} \bar{f}_T^{IR} \bar{f}_K^{TS} f_{RS}^J &= - (\lambda^2 + 1) \bar{f}_K^{IJ} \\ f_{RS}^K f_{TJ}^S \bar{f}_I^{RT} &= - (\lambda^2 + 1) f_{IJ}^K \end{aligned} \quad (4.27)$$

Now, defining the exterior product of the elements in Γ :

$$\rho \wedge \rho' \equiv \rho \otimes \rho' - \Lambda(\rho \otimes \rho') \quad (4.28)$$

we have:

$$\eta^{\hat{I}} \wedge \eta^{\hat{J}} = \left(\delta_{\hat{K}}^{\hat{I}} \delta_{\hat{L}}^{\hat{J}} - \Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}} \right) \left(\eta^{\hat{K}} \otimes \eta^{\hat{L}} \right)$$

From (4.20) and (4.22) we know that $\eta^I \wedge \eta^J$ is annihilated by the projection operator $\mathcal{P}_S$:

$$\begin{aligned} (\mathcal{P}_S)^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} \left(\eta^{\hat{K}} \wedge \eta^{\hat{L}} \right) &= 0 \\ \mathcal{P}_S &= [2]^{-2} (\Lambda + q^2) (\Lambda + q^{-2}) \\ &= [2]^{-2} \{ \Lambda + \Lambda^{-1} + (\lambda^2 + 2) \mathbf{1} \} \end{aligned} \quad (4.29)$$

namely,

$$\begin{aligned} \eta^0 \wedge \eta^0 &= 0 \\ \eta^0 \wedge \eta^I + \eta^I \wedge \eta^0 &= - \frac{\lambda}{\lambda^2 + 2} f_{JK}^I \eta^J \wedge \eta^K \\ \bar{f}_P^{IJ} f_{RS}^P \eta^R \wedge \eta^S &= - (\lambda^2 + 2) \eta^I \wedge \eta^J \end{aligned} \quad (4.30)$$

The projection operator $\mathcal{P}_A$ now can be expressed as follows:

$$\mathcal{P}_A = [2]^{-2} \{ \mathbf{2} - \Lambda - \Lambda^{-1} \} \quad (4.31)$$

It is interesting to notice that we can introduce a projection operator $\mathcal{P}_{Adj}$ with only the adjoint components that projects the product space of two adjoint representations into the adjoint representation space:

$$\begin{aligned} (\mathcal{P}_{Adj})_{KL}^{IJ} &= \frac{[2]^2}{2(\lambda^2 + 2)} (\mathcal{P}_A)_{KL}^{IJ} = - (\lambda^2 + 2)^{-1} \bar{f}_T^{IJ} f_{KL}^T \\ (\mathcal{P}_{Adj})_{RS}^{IJ} (\mathcal{P}_{Adj})_{KL}^{RS} &= (\mathcal{P}_{Adj})_{KL}^{IJ} \\ (\mathcal{P}_{Adj})_{RS}^{IJ} \bar{f}_K^{RS} &= \bar{f}_K^{IJ}, \quad (\mathcal{P}_{Adj})_{IJ}^{RS} f_{RS}^K = f_{IJ}^K \\ (\mathcal{P}_{Adj})_{KL}^{IJ} \left(\eta^K \wedge \eta^L \right) &= \eta^I \wedge \eta^J \end{aligned} \quad (4.32)$$

Now, we are in the position to define the BRST transformation δ on $\mathcal{A}$ and $\Gamma^{\otimes m}$. δ is a nilpotent operator:

$$\begin{aligned}\delta : \mathcal{A} &\rightarrow \Gamma, & \delta : \Gamma^{\wedge n} &\rightarrow \Gamma^{\wedge(n+1)} \\ \delta\alpha &= \{ig/\lambda\} (\eta^0 \alpha - \alpha \eta^0), & \alpha &\in \mathcal{A} \\ \delta\rho &= \{ig/\lambda\} \{\eta^0 \wedge \rho - (-1)^n \rho \wedge \eta^0\}, & \rho &\in \Gamma^{\wedge n}\end{aligned}\tag{4.33}$$

Introduce a functional $\chi \in \mathcal{A}'$:

$$\begin{aligned}\delta\alpha &= \eta^j (\alpha * \chi_j) \\ \chi_j &= \frac{ig}{\lambda} (\epsilon \delta_j^0 - L_j^0), & \chi_j(\mathbf{1}) &= 0 \\ \chi_j(\alpha\beta) &= \chi_j(\alpha) \epsilon(\beta) + L_j^{\hat{K}}(a) \chi_{\hat{K}}(b) \\ \chi_j \chi_{\hat{K}} &= (\chi_j \otimes \chi_{\hat{K}}) \Delta\end{aligned}\tag{4.34}$$

where χ_j are the q -analogues of the tangent vectors at the identity element of the group, and $(\cdot * \chi_j)$ are the analogues of right invariant vector fields [AC]. $\chi_{\hat{I}}(T_b^a)$ are proportional to the q -deformed Pauli matrices:

$$\begin{aligned}\chi_I(T_b^a) &= -ig [N]^{-1/2} q^{1-N+2/N} (\sigma_I)^a_b \\ \chi_0(T_b^a) &= -ig [N]^{-1/2} q^{1-N+2/N} (\sigma_0)^a_b + \{ig/\lambda\} (1 - q^{2/N}) \delta_b^a \\ &= ig \left\{ q^{-N+2/N} [N]^{-1} + \lambda^{-1} (1 - q^{2/N}) \right\} \delta_b^a\end{aligned}\tag{4.35}$$

The q -deformed structure constants can be computed from (4.34) and (4.24):

$$\begin{aligned}\mathbf{C}_{j\hat{K}}^{\hat{I}} &\equiv \chi_j(M_{\hat{K}}^{\hat{I}}), & \mathbf{C}_{j\hat{K}}^0 &= \mathbf{C}_{j0}^{\hat{K}} = 0 \\ \mathbf{C}_{0K}^J &= -ig\lambda \delta_K^J, & \mathbf{C}_{JK}^I &= -ig f_{JK}^I\end{aligned}\tag{4.36}$$

Since $\mathbf{C}_{JK}^I$ are proportional to f_{JK}^I , they satisfy the weight conservation condition:

$$\mathbf{C}_{(ij)(k\bar{\ell})}^{(r\bar{s})} = 0, \quad \text{if } \lambda_i - \lambda_{i-1} - \lambda_j + \lambda_{j-1} + \lambda_k - \lambda_{k-1} - \lambda_{\ell} + \lambda_{\ell-1} \neq \lambda_r - \lambda_{r-1} - \lambda_s + \lambda_{s-1}\tag{4.37}$$

The q -deformed Cartan-Maurer equation can be derived from (4.33):

$$\begin{aligned}
\delta\eta^0 &= \{ig/\lambda\} \{\eta^0 \wedge \eta^0 + \eta^0 \wedge \eta^0\} = 0 \\
\delta\eta^I &= \{ig/\lambda\} \{\eta^0 \wedge \eta^I + \eta^I \wedge \eta^0\} \\
&= \{ig/\lambda\} \{\eta^0 \otimes \eta^I + \eta^I \otimes \eta^0 - (\Lambda_{j\hat{K}}^{0I} + \Lambda_{j\hat{K}}^{I0}) \eta^j \otimes \eta^{\hat{K}}\} \\
&= -ig\lambda \eta^0 \otimes \eta^I - ig f_{JK}^I \eta^J \otimes \eta^K \\
&= \mathbf{C}_{j\hat{K}}^I \eta^j \otimes \eta^{\hat{K}} \\
&= \eta^j \otimes (\eta^I * \chi_j)
\end{aligned}$$

namely,

$$\begin{aligned}
\delta\eta^0 &= \mathbf{C}_{j\hat{K}}^0 \eta^j \otimes \eta^{\hat{K}} = 0 \\
\delta\eta^I &= \mathbf{C}_{j\hat{K}}^I \eta^j \otimes \eta^{\hat{K}} \\
&= \{ig/\lambda\} \{\eta^0 \wedge \eta^I + \eta^I \wedge \eta^0\} \\
&= (\lambda^2 + 2)^{-1} \mathbf{C}_{JK}^I (\eta^J \wedge \eta^K)
\end{aligned} \tag{4.38}$$

From the condition $\delta^2\alpha = 0$, the functionals χ_J span the " q -deformed Lie algebra":

$$\chi_{\hat{I}} \chi_{\hat{J}} - \Lambda_{\hat{I}\hat{J}}^{\hat{K}\hat{L}} \chi_{\hat{K}} \chi_{\hat{L}} = \mathbf{C}_{\hat{I}\hat{J}}^{\hat{K}} \chi_{\hat{K}} \tag{4.39}$$

Acting (4.39) on $M_{\hat{S}}^{\hat{P}}$, we obtain the q -deformed Jacobi identities satisfied by the q -deformed structure constants:

$$\mathbf{C}_{\hat{I}\hat{R}}^{\hat{P}} \mathbf{C}_{\hat{J}\hat{S}}^{\hat{R}} - \Lambda_{\hat{I}\hat{J}}^{\hat{K}\hat{L}} \mathbf{C}_{\hat{K}\hat{R}}^{\hat{P}} \mathbf{C}_{\hat{L}\hat{S}}^{\hat{R}} = \mathbf{C}_{\hat{I}\hat{J}}^{\hat{R}} \mathbf{C}_{\hat{R}\hat{S}}^{\hat{P}} \tag{4.40}$$

For the adjoint components we obtain from (4.40):

$$(\mathcal{P}_{Adj})_{IJ}^{KL} \mathbf{C}_{KR}^P \mathbf{C}_{LS}^R = \frac{\lambda^2 + 1}{\lambda^2 + 2} \mathbf{C}_{IJ}^R \mathbf{C}_{RS}^P \tag{4.41}$$

In fact, (4.41) is the same as the second relation in (4.27). Similarly, acting (4.39) on T_b^a we obtain following relations:

$$\chi_{\hat{I}}(T_d^a) \chi_{\hat{J}}(T_b^d) - \Lambda_{\hat{I}\hat{J}}^{\hat{K}\hat{L}} \chi_{\hat{K}}(T_d^a) \chi_{\hat{L}}(T_b^d) = \mathbf{C}_{\hat{I}\hat{J}}^{\hat{R}} \chi_{\hat{R}}(T_b^a) \tag{4.42}$$

$$(\mathcal{P}_{Adj})_{IJ}^{KL} \chi_K(T_d^a) \chi_L(T_b^d) = \xi (\lambda^2 + 2)^{-1} \mathbf{C}_{IJ}^R \chi_R(T_b^a) \tag{4.43}$$

where and hereafter, ξ denotes a constant:

$$\xi = q^{2/N} \{1 - \lambda q^{-N} [N]^{-1}\} \tag{4.44}$$

5. q -deformed BRST Algebra

Watamura [Wat] investigated the q -deformed BRST algebra $\mathcal{B}$ for $SU_q(2)$. The investigation can be generalized into the quantum groups $SU_q(N)$ straightforwardly. We sketch the main results in our notation.

$\eta^{\hat{I}}$ in the bimodule Γ is defined as the ghost field in the BRST algebra, that has the ghost number 1, but the degree of form 0. The gauge potential $A^{\hat{I}}$ has the degree of form 1, but the ghost number 0. There are two nilpotent operators in the BRST algebra: The operator δ increases the ghost number by one, and the operator d increases the degree of form by one. Neglecting the matter field, that is irrelevant to our following discussion, we are only interested in four fields in the BRST algebra $\mathcal{B}$: η , $d\eta$, A , and dA , which satisfy the following algebraic relations.

Firstly, we introduce an index n that is equal to the difference between the degree of form and the ghost number. The indices n for η , $d\eta$, A and dA are -1, 0, 1, and 2, respectively. Both nilpotent operators δ and d satisfy the Leibniz rule in the graded sense for the index n :

$$\begin{aligned}\delta^2 &= 0, & d^2 &= 0, & d\delta + \delta d &= 0 \\ d(XY) &= (dX)Y + (-1)^{n_x} X(dY) \\ \delta(XY) &= (\delta X)Y + (-1)^{n_x} X(\delta Y)\end{aligned}\tag{5.1}$$

where $X, Y \in \mathcal{B}$, and n_x is the index of X . Both d and δ are covariant for the left and right actions: For any element $X \in \mathcal{B}$ they satisfy:

$$\begin{aligned}\Delta_L(\delta X) &= (id \otimes \delta)\Delta_L(X), & \Delta_L(dX) &= (id \otimes d)\Delta_L(X) \\ \Delta_R(\delta X) &= (\delta \otimes id)\Delta_R(X), & \Delta_R(dX) &= (d \otimes id)\Delta_R(X)\end{aligned}\tag{5.2}$$

Secondly, the gauge potentials $A^{\hat{I}}$ are assumed [Wat] to have similar properties like $\eta^{\hat{I}}$. Hereafter, we neglect the wedge sign $\wedge$ for simplicity.

$$\begin{aligned}(\mathcal{P}_S)^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} (A^{\hat{K}} A^{\hat{L}}) &= 0 \\ A^0 A^0 &= 0, & (\mathcal{P}_{Adj})^{IJ}_{KL} (A^K A^L) &= A^I A^J \\ \{ig/\lambda\} (A^0 A^I + A^I A^0) &= (\lambda^2 + 2)^{-1} \mathbf{C}_{JK}^I A^J A^K\end{aligned}\tag{5.3}$$

From the consistent conditions [Wat], $d\eta^{\hat{J}}$ and $dA^{\hat{J}}$ have to satisfy another relation:

$$(\mathcal{P}_A)^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} (d\eta^{\hat{K}} d\eta^{\hat{L}}) = 0, \quad (\mathcal{P}_A)^{\hat{I}\hat{J}}_{\hat{K}\hat{L}} (dA^{\hat{K}} dA^{\hat{L}}) = 0\tag{5.4}$$

namely,

$$\begin{aligned} d\eta^I d\eta^0 &= d\eta^0 d\eta^I = -\lambda^{-1} f_{JK}^I d\eta^J d\eta^K \\ dA^I dA^0 &= dA^0 dA^I = -\lambda^{-1} f_{JK}^I dA^J dA^K \end{aligned}$$

Thirdly, the gauge potential is introduced in the covariant derivative. The covariant condition of the covariant derivative in the BRST transformation requires:

$$\delta A^0 = 0, \quad \delta A^I = d\eta^I + \frac{ig}{\lambda} (\eta^0 A^I + A^I \eta^0) \quad (5.5)$$

Fourthly, for two different fields $X^{\hat{J}}$ and $Y^{\hat{K}}$ in $\mathcal{B}$ with indices n_x and n_y , $n_x > n_y$, respectively, the consistent condition requires the following commutative relations:

$$(-1)^{n_x n_y} X^{\hat{I}} Y^{\hat{J}} = Y^{\hat{K}} (X^{\hat{I}} * L_{\hat{K}}^{\hat{J}}) = \Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}} Y^{\hat{K}} X^{\hat{L}} \quad (5.6)$$

From (4.24) we have

$$\begin{aligned} (-1)^{n_x n_y} X^0 Y^{\hat{J}} &= Y^{\hat{J}} X^0 \\ \{ig/\lambda\} (Y^0 X^I - (-1)^{n_x n_y} X^I Y^0) & \\ = -ig\lambda Y^0 X^I - ig f_{JK}^I Y^J X^K &= Y^{\hat{J}} X^{\hat{K}} \mathbf{C}_{\hat{J}\hat{K}}^I \end{aligned} \quad (5.7)$$

At last, the gauge fields F^J satisfy:

$$\begin{aligned} F^{\hat{J}} &= dA^{\hat{J}} + \{ig/\lambda\} (A^0 A^{\hat{J}} + A^{\hat{J}} A^0) \\ F^0 &= dA^0, \quad F^I = dA^I + (\lambda^2 + 2)^{-1} \mathbf{C}_{JK}^I A^J A^K \\ F^{\hat{I}} \eta^{\hat{J}} &= \eta^{\hat{K}} F^{\hat{L}} \Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}} \end{aligned} \quad (5.8)$$

$$\delta F^{\hat{I}} = \{ig/\lambda\} (\eta^0 F^{\hat{I}} - F^{\hat{I}} \eta^0) = \eta^{\hat{J}} F^{\hat{K}} \mathbf{C}_{\hat{J}\hat{K}}^{\hat{I}} \quad (5.9)$$

$$\begin{aligned} dF^{\hat{I}} &= -\{ig/\lambda\} (A^0 F^{\hat{I}} - F^{\hat{I}} A^0) \\ &= -\{ig/\lambda\} (A^0 dA^{\hat{I}} - dA^{\hat{I}} A^0) \\ &= -A^{\hat{J}} dA^{\hat{K}} \mathbf{C}_{\hat{J}\hat{K}}^{\hat{I}} \end{aligned} \quad (5.10)$$

The commutative relation (5.6) can be rewritten as follows.

Proposition 2. *The components X_b^a and Y_b^a of two different fields $X^{\hat{J}}$ and $Y^{\hat{K}}$ in $\mathcal{B}$ with indices n_x and n_y , $n_x > n_y$, respectively, satisfy:*

i)

$$(-1)^{n_x n_y} X^a_r \left(\hat{R}_q^{-1} \right)^{ri}_{sk} Y^s_t \left(\hat{R}_q^{-1} \right)^{tk}_{bj} = \left(\hat{R}_q^{-1} \right)^{ai}_{rk} Y^r_s \left(\hat{R}_q^{-1} \right)^{sk}_{tj} X^t_b \quad (5.11)$$

ii)

$$\begin{aligned} & \{ig/\lambda\} \left(Y^0 X^I - (-1)^{n_x n_y} X^I Y^0 \right) (\sigma_I)^a_b \\ &= -ig q^{1-N} [N]^{-1/2} \left(Y^a_d X^d_b - (-1)^{n_x n_y} X^a_d Y^d_b \right) \end{aligned} \quad (5.12)$$

where

$$X^a_b = X^{\hat{I}} (\sigma_{\hat{I}})^a_b, \quad Y^a_b = Y^{\hat{I}} (\sigma_{\hat{I}})^a_b$$

Proof. (5.6) can be rewritten in terms of the explicit form (4.17) of $\Lambda^{\hat{I}\hat{J}}_{\hat{K}\hat{L}}$:

$$\begin{aligned} & (-1)^{n_x n_y} (\sigma_{\hat{I}})^{a\bar{b}} (\sigma_{\hat{J}})^{c\bar{d}} X^{\hat{I}} Y^{\hat{J}} \\ &= \left(\hat{R}_q^{-1} \right)^{ar}_{iu} \left(\hat{R}_q \right)^{u\bar{s}}_{\bar{j}k} \left(\hat{R}_q^{-1} \right)^{\bar{b}c}_{r\bar{v}} \left(\hat{R}_q \right)^{\bar{v}\bar{d}}_{\bar{s}\bar{\ell}} (\sigma_{\hat{K}})^{i\bar{j}} (\sigma_{\hat{L}})^{k\bar{\ell}} Y^{\hat{K}} X^{\hat{L}} \end{aligned} \quad (5.13)$$

Moving two factors $\left(\hat{R}_q^{-1} \right)^{\bar{b}c}_{r\bar{v}} \left(\hat{R}_q \right)^{\bar{v}\bar{d}}_{\bar{s}\bar{\ell}}$ from the right hand side of (5.13) to the left, and left multiplying (5.13) by $q^{-1} \epsilon_{\bar{s}s'} \epsilon_{\bar{\ell}\ell'}$, we obtain the left hand side of the equation as follows:

$$\begin{aligned} & (-1)^{n_x n_y} q^{-1} \epsilon_{\bar{s}s'} \epsilon_{\bar{\ell}\ell'} \left(\hat{R}_q^{-1} \right)^{\bar{s}\bar{\ell}}_{\bar{v}\bar{d}} \left(\hat{R}_q \right)^{r\bar{v}}_{\bar{b}c} (\sigma_{\hat{I}})^{a\bar{b}} (\sigma_{\hat{K}})^{c\bar{d}} X^{\hat{I}} Y^{\hat{J}} \\ &= (-1)^{n_x n_y} q^{-1} \epsilon_{\bar{s}s'} \left(\hat{R}_q^{-1} \right)^{br}_{cv} \left(\hat{R}_q \right)^{v\bar{s}}_{\bar{d}\ell'} X^a_b Y^c_{d'} \epsilon^{d'\bar{d}} \\ &= (-1)^{n_x n_y} X^a_b \left(\hat{R}_q^{-1} \right)^{br}_{cv} Y^c_d \left(\hat{R}_q^{-1} \right)^{d\bar{v}}_{\ell's'} \end{aligned}$$

The right hand side becomes:

$$\begin{aligned} & q^{-1} \epsilon_{\bar{s}s'} \epsilon_{\bar{\ell}\ell'} \left(\hat{R}_q^{-1} \right)^{ar}_{iu} \left(\hat{R}_q \right)^{u\bar{s}}_{\bar{j}k} (\sigma_{\hat{K}})^{i\bar{j}} (\sigma_{\hat{L}})^{k\bar{\ell}} Y^{\hat{K}} X^{\hat{L}} \\ &= \left(\hat{R}_q^{-1} \right)^{ar}_{iu} Y^i_j \left(\hat{R}_q^{-1} \right)^{ju}_{ks'} X^k_{\ell'} \end{aligned}$$

Comparing two sides of (5.13) we obtain (5.11). From (5.7) and (4.42) we have:

$$\begin{aligned} & \{ig/\lambda\} \left(Y^0 X^I - (-1)^{n_x n_y} X^I Y^0 \right) (\sigma_I)^a_b \\ &= (\sigma_I)^a_b \mathbf{C}^I_{j\hat{K}} Y^j X^{\hat{K}} \\ &= \left(-igq^{1-N+2/N} [N]^{-1/2} \right)^{-1} \chi_{\hat{I}}(T^a_b) \mathbf{C}^{\hat{I}}_{j\hat{K}} Y^j X^{\hat{K}} \\ &= \left(-igq^{1-N+2/N} [N]^{-1/2} \right)^{-1} \left\{ Y^{\hat{J}} \chi_{\hat{J}}(T^a_d) X^{\hat{K}} \chi_{\hat{K}}(T^d_b) \right. \\ &\quad \left. - (-1)^{n_x n_y} X^{\hat{J}} \chi_{\hat{J}}(T^a_d) Y^{\hat{K}} \chi_{\hat{K}}(T^d_b) \right\} \\ &= \left(-igq^{1-N+2/N} [N]^{-1/2} \right) \left(Y^a_d X^d_b - (-1)^{n_x n_y} X^a_d Y^d_b \right) \\ &\quad + \{ig/\lambda\} (1 - q^{2/N}) (Y^0 X^a_b - (-1)^{n_x n_y} X^a_b Y^0) \\ &= \left(-igq^{1-N} [N]^{-1/2} \right) \left(Y^a_d X^d_b - (-1)^{n_x n_y} X^a_d Y^d_b \right) \end{aligned}$$

(5.12) was proved. Q.E.D.

6. q -Deformed Chern Class and q -Trace

In our previous paper [HHM], omitting the possible constant factor, we assume that the second q -deformed Chern class P for the quantum group $SU_q(2)$ has the following form:

$$P = \langle F, F \rangle \equiv F^I F^J g'_{IJ} \quad (6.1)$$

where from the condition:

$$\delta P = 0, \quad dP = 0 \quad (6.2)$$

we defined the q -deformed Killing form g'_{IJ} as:

$$g'_{IJ} = D^R_S \mathbf{C}_{IT}^S \mathbf{C}_{JR}^T \quad (6.3)$$

For $SU_q(N)$ the non-vanishing components are:

$$\begin{aligned} g'_{(k\bar{k})(k\bar{k})} &= -g^2 q^{2k-2N} \{[2N]/[N]\} \\ g'_{(j\bar{k})(k\bar{j})} &= -g^2 q^{j-k-N+2} \{[2N]/[N]\}, \quad j \neq k \end{aligned}$$

where the repeated indices are not summed.

Recalling (4.35) and (4.36), we may define another q -deformed Killing form g_{IJ} :

$$\begin{aligned} g_{IJ} &= D^a_b \chi_I(T^b_c) \chi_J(T^c_a) \\ g_{(k\bar{k})(k\bar{k})} &= -g^2 [N] q^{2k-3N+4/N} \\ g_{(j\bar{k})(k\bar{j})} &= -g^2 [N] q^{j-k+2-2N+4/N}, \quad j \neq k \end{aligned} \quad (6.4)$$

where the repeated indices are not summed. The else components of g_{JK} are vanishing. Both for $SU_q(2)$ and for $SU_q(N)$ two q -deformed Killing forms are proportional to each other, namely, just like the Killing form in a Lie algebra, the q -deformed Killing form is also independent of the representation in which it is calculated.

Now, we are going to define the higher q -deformed Chern class P_m for the quantum group $SU_q(N)$ from the covariant condition (6.2). Generalizing (6.4) we define the "generalized q -deformed Killing forms" and the q -trace as follows:

$$g_{I_1 I_2 \dots I_m} = D^{a_0}_{a_1} \chi_{I_1}(T^{a_1}_{a_2}) \chi_{I_2}(T^{a_2}_{a_3}) \dots \chi_{I_m}(T^{a_m}_{a_0}) \quad (6.5)$$

$$\langle X_1, X_2, \dots, X_m \rangle = X_1^{I_1} X_2^{I_2} \dots X_m^{I_m} g_{I_1 I_2 \dots I_m} \quad (6.6)$$

where X_i are fields η , $d\eta$, A or dA in the BRST algebra $\mathcal{B}$. In (6.6) the fields can also be replaced by, for example, XY^0 , Y^0X or F . From the properties of q -Pauli matrices, the sum of all subscripts of nonvanishing components $g_{I_1 \dots I_m}$, as weights, has to be zero. The following theorem is easily proved from (4.43) and the definition (6.5).

Proposition 3. *The generalized q -deformed Killing forms satisfy the following relations:*

$$(\mathcal{P}_{Adj})_{JK}^{RS} g_{I_1 \dots I_{n-1} R S I_{n+2} \dots I_m} = -ig\xi (\lambda^2 + 2)^{-1} f_{JK}^R g_{I_1 \dots I_{n-1} R I_{n+2} \dots I_m} \quad (6.7)$$

where ξ was given in (4.44).

(6.7) can be rewritten in another form by removing a factor f_{JK}^P :

$$\bar{f}_K^{RS} g_{I_1 \dots I_{n-1} R S I_{n+2} \dots I_m} = ig \xi g_{I_1 \dots I_{n-1} K I_{n+2} \dots I_m} \quad (6.8)$$

The m -th q -deformed Chern class P_m for the quantum group $SU_q(N)$ is defined as follows:

$$\begin{aligned} P_m &= \langle F, F, \dots, F \rangle \\ &= F^{I_1} F^{I_2} \dots F^{I_m} g_{I_1 I_2 \dots I_m} \\ &= \left(-ig[N]^{-1/2} q^{1-N+2/N} \right)^m D_{a_1}^{a_0} F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_0}^{a_m} \end{aligned} \quad (6.9)$$

From (5.9) and (5.12) we have:

$$\begin{aligned} \delta P_m &= q^{2m/N} \left(-igq^{1-N}[N]^{-1/2} \right)^{m+1} \left\{ D_{a_1}^{a_0} \eta_{a_1}^b F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_0}^{a_m} \right. \\ &\quad \left. - D_{a_1}^b F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_0}^{a_m} \eta_{a_1}^{a_0} \right\} \end{aligned} \quad (6.10)$$

By (2.14), (2.15) and (5.11) the second term cancels the first term:

$$\begin{aligned} &D_{a_1}^b F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_0}^{a_m} \eta_{a_1}^{a_0} \\ &= D_{a_1}^b F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_0}^{a_m} \left\{ D_{a_1}^i q^N \left(\hat{R}_q^{-1} \right)_{di}^{a_0 j} \right\} \eta_{a_1}^d \\ &= q^N D_{a_1}^b D_{a_1}^i F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_m}^{a_{m-1}} F_{a_0}^{a_m} \left(\hat{R}_q^{-1} \right)_{dk}^{a_0 j} \eta_c^d \left(\hat{R}_q^{-1} \right)_{r\ell}^{ck} \left(\hat{R}_q \right)_{bi}^{r\ell} \\ &= q^N D_{a_1}^b D_{a_1}^i F_{a_2}^{a_1} F_{a_3}^{a_2} \dots F_{a_m}^{a_{m-1}} \left(\hat{R}_q^{-1} \right)_{dk}^{a_m j} \eta_c^d \left(\hat{R}_q^{-1} \right)_{r\ell}^{ck} F_{a_0}^r \left(\hat{R}_q \right)_{bi}^{a_0 \ell} \\ &= q^N D_{a_1}^b D_{a_1}^i \left(\hat{R}_q^{-1} \right)_{dk}^{a_1 j} \eta_c^d \left(\hat{R}_q^{-1} \right)_{r\ell}^{ck} F_{a_2}^r F_{a_3}^{a_2} \dots F_{a_m}^{a_{m-1}} F_{a_0}^{a_m} \left(\hat{R}_q \right)_{bi}^{a_0 \ell} \\ &= q^N D_{a_1}^{a_0} D_{a_1}^i \eta_c^{a_1} \left(\hat{R}_q^{-1} \right)_{ri}^{cj} F_{a_2}^r F_{a_3}^{a_2} \dots F_{a_m}^{a_{m-1}} F_{a_0}^{a_m} \\ &= D_{a_1}^{a_0} \eta_{a_1}^r F_{a_2}^r F_{a_3}^{a_2} \dots F_{a_m}^{a_{m-1}} F_{a_0}^{a_m} \end{aligned}$$

Thus, $\delta P_m = 0$. This technique of proof was firstly used by Isaev [Isa]. The proof of $dP_m = 0$ can be performed analogously:

$$\begin{aligned} dP_m &= \left(gq^{1-N+1/N} [N]^{-1/2} \right)^{2m} \left\{ D_{b}^{a_0} A_{a_1}^b dA_{a_2}^{a_1} dA_{a_3}^{a_2} \cdots dA_{a_0}^{a_m} \right. \\ &\quad \left. - D_{a_1}^b dA_{a_2}^{a_1} dA_{a_3}^{a_2} \cdots dA_{a_0}^{a_m} A_{b}^{a_0} \right\} \\ &= 0 \end{aligned}$$

Note that the components of the identity and the adjoint representations are separated in the q -deformed Chern class, although they are mixed in the commutative relations of BRST algebra.

This technique can be used to prove more general relations. Remind (6.6) and that the condition $\delta P_m = 0$ can be rewritten as follows:

$$\begin{aligned} \delta P_m &= \{ig/\lambda\} \left\{ \eta^0 F^{I_1} F^{I_2} \cdots F^{I_m} g_{I_1 I_2 \cdots I_m} - F^{I_1} F^{I_2} \cdots F^{I_m} \eta^0 g_{I_1 I_2 \cdots I_m} \right\} \\ &= 0 \end{aligned}$$

Now, the following theorem can be proved straightforwardly.

Proposition 4. *Let $Y \in \mathcal{B}$ with index n_y , and X_i , that are not the field Y , be any fields in $\mathcal{B}$ with the indices n_i , $n_i > n_y$. Then:*

$$\begin{aligned} (-1)^n \langle X_1, \cdots, X_m, Y \rangle &= \langle Y, X_1, \cdots, X_m \rangle \\ (-1)^n \langle X_1, \cdots, X_m \rangle Y^0 &= Y^0 \langle X_1, \cdots, X_m \rangle \\ n &= \sum_{i=1}^m n_i n_y \end{aligned} \tag{6.11}$$

Some corollaries can be derived from Proposition 4. First of all, substituting the commutative relation (5.6) into (6.11) we obtain some constraints for the "generalized q -deformed Killing forms":

$$\begin{aligned} g_{I_1 I_2 \cdots I_m I_0} \Lambda_{J_0 J_1}^{I_1 \hat{K}_1} \Lambda_{J_1 J_2}^{I_2 \hat{K}_2} \cdots \Lambda_{J_{m-2} J_{m-1}}^{I_{m-1} \hat{K}_{m-1}} \Lambda_{J_{m-1} J_m}^{I_m I_0} &= g_{J_0 J_1 \cdots J_m} \\ g_{I_1 I_2 \cdots I_m I_0} \Lambda_{0 J_1}^{I_1 \hat{K}_1} \Lambda_{J_1 J_2}^{I_2 \hat{K}_2} \cdots \Lambda_{J_{m-2} J_{m-1}}^{I_{m-1} \hat{K}_{m-1}} \Lambda_{J_{m-1} J_m}^{I_m I_0} &= 0 \end{aligned} \tag{6.12}$$

$$g_{I_1 I_2 \cdots I_m} \Lambda_{J_1}^{I_1 \hat{K}_1} \Lambda_{J_2}^{I_2 \hat{K}_2} \cdots \Lambda_{J_{m-2} J_{m-1}}^{I_{m-1} \hat{K}_{m-1}} \Lambda_{J_{m-1} J_m}^{I_m 0} = \delta_{\hat{K}_0}^0 g_{J_1 J_2 \cdots J_m} \tag{6.13}$$

(6.12) describes the cyclic property of the q -trace. (6.13) is equivalent to the following form:

$$\sum_{n=1}^m g_{I_1 \dots I_n J_{n+1} \dots J_m} \Lambda_{\hat{K}_0 J_1}^{I_1 \hat{K}_1} \Lambda_{\hat{K}_1 J_2}^{I_2 \hat{K}_2} \dots \Lambda_{\hat{K}_{n-2} J_{n-1}}^{I_{n-1} \hat{K}_{n-1}} \mathbf{C}_{\hat{K}_{n-1} J_n}^{I_n} = 0 \quad (6.14)$$

namely,

$$g_{I_1 I_2 \dots I_m} \chi_{\hat{K}_0} \left(M_{J_1}^{I_1} \dots M_{J_m}^{I_m} \right) = 0 \quad (6.15)$$

Bernard [Ber] gave the special form ($m = 2$) of (6.14):

$$g_{I_1 J_2} \mathbf{C}_{\hat{K}_0 \hat{J}_1}^{I_1} + g_{I_1 I_2} \Lambda_{\hat{K}_0 \hat{J}_1}^{I_1 \hat{K}_1} \mathbf{C}_{\hat{K}_1 \hat{J}_2}^{I_2} = 0 \quad (6.16)$$

In terms of Proposition 3, it can be proved by direct calculation that the constraint (6.12) is equivalent to (6.13). It is not surprising because they come from the same source. In fact, by making use of the explicit form (4.24) we obtain the same relations from (6.13) as from (6.12). For example, when $m = 1$ we obtain:

$$g_I = 0 \quad (6.17)$$

It implies the orthogonality (3.9) of q -Pauli matrices. When $m = 2$, we obtain:

$$\begin{aligned} (\lambda^2 + 2) g_{IJ} + g_{RS} \bar{f}_I^{RT} f_{TJ}^S &= 0 \\ (\lambda^2 + 1) g_{RK} f_{IJ}^R + g_{IR} f_{JK}^R + g_{RS} \bar{f}_P^{RT} f_{IJ}^P f_{TK}^S &= 0 \end{aligned}$$

Thus,

$$f_{IJ}^R g_{RK} = g_{IR} f_{JK}^R \quad (6.18)$$

When $m = 3$ we have:

$$\begin{aligned} &(\lambda^4 + 3\lambda^2 + 3) g_{IJK} + (\lambda^2 + 1) \left\{ \bar{f}_I^{RT} f_{TJ}^S g_{RSK} + \bar{f}_J^{RT} f_{TK}^S g_{IRS} \right\} \\ &+ \bar{f}_I^{RS} f_{JK}^T g_{RST} + \bar{f}_I^{RP_3} f_{P_3 J}^{P_2} \bar{f}_{P_2}^{SP_1} f_{P_1 K}^T g_{RST} = 0 \\ &(\lambda^2 + 2) f_{LI}^R g_{RJK} - (\lambda^2 + 1) f_{IJ}^R g_{LRK} - f_{JK}^R g_{LIR} \\ &= \bar{f}_P^{RT} f_{IJ}^P f_{TK}^S g_{LRS} - f_{LI}^R \bar{f}_J^{SP} f_{PK}^T g_{RST} \end{aligned} \quad (6.19)$$

At last, from Propositions 3 and 4 we can prove:

$$\langle A^m \rangle \equiv A^{I_1} A^{I_2} \dots A^{I_m} g_{I_1 I_2 \dots I_m} = 0, \quad \text{when } m \neq 3 \quad (6.20)$$

In fact, define:

$$\begin{aligned}
\langle A^m \rangle \phi_n &= g_{T_1 \dots T_{m-n} I_1 \dots I_n} \Lambda_{J_1 J_2}^{I_1 \hat{K}_1} \Lambda_{\hat{K}_1 J_3}^{I_2 \hat{K}_2} \dots \Lambda_{\hat{K}_{n-3} J_{n-1}}^{I_{n-2} \hat{K}_{n-2}} \Lambda_{\hat{K}_{n-2} J_n}^{I_{n-1} I_n} \\
&\quad \cdot A^{T_1} \dots A^{T_{m-n}} A^{J_1} \dots A^{J_n}, \quad 2 \leq n \leq m \\
\langle A^m \rangle \psi_n &= g_{T_1 \dots T_{m-n} I_1 \dots I_n} \Lambda_{J_1 J_2}^{I_1 0} \Lambda_{0 J_3}^{I_2 \hat{K}_2} \dots \Lambda_{\hat{K}_{n-3} J_{n-1}}^{I_{n-2} \hat{K}_{n-2}} \Lambda_{\hat{K}_{n-2} J_n}^{I_{n-1} I_n} \\
&\quad \cdot A^{T_1} \dots A^{T_{m-n}} A^{J_1} \dots A^{J_n}, \quad 3 \leq n \leq m
\end{aligned} \tag{6.21}$$

Since the factor $\langle A^m \rangle$ has been abstracted, the functions ϕ_n and ψ_n are independent of m . From (4.24), (4.27) and (5.3) we obtain the recurrence relations and some explicit values for the functions:

$$\begin{aligned}
\phi_n &= -(\lambda^2 + 1) \phi_{n-1} + \psi_n, \quad n \geq 3 \\
\psi_n &= (\lambda^2 + 1) \psi_{n-1} - \lambda^2 (\lambda^2 + 1) \psi_{n-2} \\
&\quad + \sum_{a=1}^{n-5} \left\{ (-1)^{a+1} \lambda^2 \left((\lambda^2 + 1)^2 - a(\lambda^2 + 2) \right) \psi_{n-a-2} \right\} \\
&\quad + (-1)^n \lambda^2 (\lambda^2 + 2) (\lambda^2 + 1)^{n-4} (\lambda^2 - n + 4), \quad n \geq 5 \\
\phi_2 &= -\lambda^2 - 1, \quad \phi_3 = 1, \quad \phi_4 = -\lambda^2 (\lambda^2 + 3) - 1 \\
\phi_5 &= \lambda^2 (\lambda^2 + 2)^2 + 1, \quad \phi_6 = -\lambda^2 (\lambda^2 + 3) (\lambda^4 + 3\lambda^2 + 3) - 1
\end{aligned}$$

The limit values of the functions are as follows:

$$\begin{aligned}
\lim_{q \rightarrow 1} \phi_{2n} &= -1, \quad \lim_{q \rightarrow 1} \phi_{2n+1} = 1, \quad \lim_{q \rightarrow 1} \psi_n = 0 \\
\lim_{q \rightarrow 1} \lambda^{-2} (\phi_{2n} + 1) &= -2n^2 + 4n - 3 \\
\lim_{q \rightarrow 1} \lambda^{-2} (\phi_{4n+1} - 1) &= 4n(2n - 1), \quad \lim_{q \rightarrow 1} \lambda^{-2} (\phi_{4n+3} + 1) = 4n(2n + 1) \\
\lim_{q \rightarrow 1} \lambda^{-2} \psi_{2n} &= -2(n - 1), \quad \lim_{q \rightarrow 1} \lambda^{-2} \psi_{2n+1} = 2(n - 2)
\end{aligned}$$

Now, substituting (6.12) into (6.21) we have:

$$(\phi_m - 1) \langle A^m \rangle = 0 \tag{6.22}$$

It leads to (6.20). It is interesting to notice that in the classical case ($q = 1$) only $\langle A^{2n} \rangle = 0$ is well known. (6.20) also holds for η instead of A due to (4.32).

Generalize the q -trace (6.6) as follows:

$$\langle \dots, Z_1, [X, Y], Z_2, \dots \rangle = \dots Z_1^{I_1} X^J Y^K Z_2^{I_2} \dots (\mathcal{P}_{Adj})_{JK}^{RS} g_{\dots I_1 R S I_2 \dots} \tag{6.23}$$

where X, Y and Z are fields in the BRST algebra $\mathcal{B}$. The fields can also be replaced by, for example, XY^0, Y^0X or F . From Propositions 3 and 4, we have:

$$\begin{aligned}
\langle \cdots, X, [\eta, \eta], Y, \cdots \rangle &= \langle \cdots, X, \eta, \eta, Y, \cdots \rangle \\
\langle \cdots, X, [A, A], Y, \cdots \rangle &= \langle \cdots, X, A, A, Y, \cdots \rangle \\
\langle \cdots, Z_1, [X, Y], Z_2, \cdots \rangle & \\
&= \cdots Z_1^{I_1} \left\{ -ig\xi(\lambda^2 + 2)^{-1} f_{JK}^T X^J Y^K \right\} Z_2^{I_2} \cdots g_{\cdots I_1 T I_2 \cdots}
\end{aligned} \tag{6.24}$$

$$(\mathcal{P}_{Adj})_{JK}^{RS} g_{RS} = 0, \quad \bar{f}_J^{RS} g_{RS} = 0, \quad \Lambda_{JK}^{RS} g_{RS} = g_{JK} \tag{6.25}$$

$$(\mathcal{P}_{Adj})_{IJ}^{RS} g_{RSK} = g_{IRS} (\mathcal{P}_{Adj})_{JK}^{RS}, \quad \langle X, [Y, Z] \rangle = \langle [X, Y], Z \rangle \tag{6.26}$$

where X, Y and Z are fields in the BRST algebra $\mathcal{B}$.

Proposition 5. Assume X^J and Y^J are two different fields in $\mathcal{B}$ with the indices $n_x > n_y$. Z^J denotes a field in $\mathcal{B}$, or a field multiplying by the zero component of this or another field. Since the following relations are linear for the field Z^J , Z^J can also be replaced by their linear combination, for example, F^J . Then,

$$\begin{aligned}
(-1)^{n_x n_y} \langle \cdots, Z_1, XY^0, Z_2, \cdots \rangle & \\
= (\lambda^2 + 1) \langle \cdots, Z_1, Y^0 X, Z_2, \cdots \rangle - \frac{\lambda(\lambda^2 + 2)}{ig\xi} \langle \cdots, Z_1, [Y, X], Z_2, \cdots \rangle
\end{aligned} \tag{6.27}$$

$$\begin{aligned}
\{ig/\lambda\} \langle \cdots, Z_1, (A^0 A + A A^0), Z_2, \cdots \rangle & \\
= \xi^{-1} \langle \cdots, Z_1, A, A, Z_2, \cdots \rangle & \\
\{ig/\lambda\} \langle \cdots, Z_1, (\eta^0 \eta + \eta \eta^0), Z_2, \cdots \rangle & \\
= \xi^{-1} \langle \cdots, Z_1, \eta, \eta, Z_2, \cdots \rangle &
\end{aligned} \tag{6.28}$$

$$\begin{aligned}
(-1)^{n_x n_y} \langle \cdots, Z_1, X, Y, Z_2, \cdots \rangle & \\
= \langle \cdots, Z_1, Y, X, Z_2, \cdots \rangle & \\
- ig\lambda^{-1} \xi \langle \cdots, Z_1, (Y^0 X - (-1)^{n_x n_y} XY^0), Z_2, \cdots \rangle &
\end{aligned} \tag{6.29}$$

$$\begin{aligned}
& (-1)^{n_x n_y} \langle \cdots, Z_1, X, Y, Z_2, \cdots \rangle \\
&= ig\lambda\xi \langle \cdots, Z_1, Y^0 X, Z_2, \cdots \rangle \\
&\quad - (\lambda^2 + 2) \langle \cdots, Z_1, [Y, X], Z_2, \cdots \rangle + \langle \cdots, Z_1, Y, X, Z_2, \cdots \rangle \\
& (-1)^{n_x n_y} \langle \cdots, Z_1, [X, Y], Z_2, \cdots \rangle \\
&= ig\lambda\xi \langle \cdots, Z_1, Y^0 X, Z_2, \cdots \rangle - (\lambda^2 + 1) \langle \cdots, Z_1, [Y, X], Z_2, \cdots \rangle
\end{aligned} \tag{6.30}$$

Proof. From (5.7) and (4.36) we have

$$(-1)^{n_x n_y} X^I Y^0 = (\lambda^2 + 1) Y^0 X^I + \lambda Y^J X^K f_{JK}^I$$

Due to (4.43),

$$\begin{aligned}
(-1)^{n_x n_y} X^I Y^0 \chi_I(T_b^a) &= (\lambda^2 + 1) Y^0 X^I \chi_I(T_b^a) \\
&\quad - \frac{\lambda(\lambda^2 + 2)}{ig\xi} Y^I X^L (\mathcal{P}_{Adj})_{IL}^{JK} \chi_J(T_d^a) \chi_K(T_b^d)
\end{aligned}$$

According to the definition (6.6) we obtain (6.27).

From (5.3) we have:

$$\begin{aligned}
\{ig/\lambda\} (A^0 A^I + A^I A^0) \chi_I(T_b^a) &= (\lambda^2 + 2)^{-1} A^J A^K \mathbf{C}_{JK}^I \chi_I(T_b^a) \\
&= \xi^{-1} A^J A^K \chi_J(T_d^a) \chi_K(T_b^d)
\end{aligned}$$

This relation leads to (6.28). The relation also holds for η^J due to (4.32).

From (4.42) and (5.7) we obtain:

$$\begin{aligned}
(-1)^{n_x n_y} X^{\hat{I}} Y^{\hat{J}} \chi_{\hat{I}}(T_d^a) \chi_{\hat{J}}(T_b^d) &= Y^{\hat{K}} X^{\hat{L}} \Lambda_{\hat{K}\hat{L}}^{\hat{I}\hat{J}} \chi_{\hat{I}}(T_d^a) \chi_{\hat{J}}(T_b^d) \\
&= Y^{\hat{K}} X^{\hat{L}} \left\{ \chi_{\hat{K}}(T_d^a) \chi_{\hat{L}}(T_b^d) - \mathbf{C}_{\hat{K}\hat{L}}^T \chi_T(T_b^a) \right\} \\
&= Y^{\hat{K}} X^{\hat{L}} \chi_{\hat{K}}(T_d^a) \chi_{\hat{L}}(T_b^d) - \{ig/\lambda\} (Y^0 X^J - (-1)^{n_x n_y} X^J Y^0) \chi_J(T_b^a)
\end{aligned}$$

Then, noting (4.35) and (5.7) we have:

$$\begin{aligned}
& \{(-1)^{n_x n_y} X^I Y^J - Y^I X^J\} \chi_I(T_d^a) \chi_J(T_b^d) \\
&= \{(-1)^{n_x n_y} X^{\hat{I}} Y^{\hat{J}} - Y^{\hat{I}} X^{\hat{J}}\} \chi_{\hat{I}}(T_d^a) \chi_{\hat{J}}(T_b^d) \\
&\quad - ig \left\{ q^{-N+2/N} [N]^{-1} + \lambda^{-1} (1 - q^{2/N}) \right\} \{(-1)^{n_x n_y} X^J Y^0 - Y^0 X^J\} \chi_J(T_b^a) \\
&= - ig\lambda^{-1}\xi \{Y^0 X^J - (-1)^{n_x n_y} X^J Y^0\} \chi_J(T_b^a)
\end{aligned}$$

It leads to (6.29). From (6.27) and (6.29) we obtain (6.30). Q.E.D.

From Proposition 5 we obtain:

$$\begin{aligned}\langle F, A \rangle &= \langle A, F \rangle = \langle A, dA \rangle + \{ig/\lambda\} \langle A, (A^0 A + A A^0) \rangle \\ &= \langle A, dA \rangle + \xi^{-1} \langle A, A, A \rangle\end{aligned}\quad (6.31)$$

Proposition 6. *For any field $X \in \mathcal{B}$,*

$$\langle \eta, \eta, [\eta, X] \rangle = 0, \quad \langle A, A, [A, X] \rangle = 0 \quad (6.32)$$

Proof. From (4.27) and (4.32) we have:

$$\begin{aligned}f_{IL}^T f_{JK}^L \eta^I \eta^J X^K &= -(\lambda^2 + 2)^{-1} f_{IL}^T f_{JK}^L \bar{f}_P^{IJ} f_{RS}^P \eta^R \eta^S X^K \\ &= \frac{\lambda^2 + 1}{\lambda^2 + 2} f_{RS}^P f_{PK}^T \eta^R \eta^S X^K\end{aligned}\quad (6.33)$$

Noting (6.7) and (6.18) we have:

$$\begin{aligned}\left(\frac{\lambda^2 + 2}{-ig\xi}\right)^2 \langle \eta, \eta, [\eta, X] \rangle &= \eta^I \eta^J \eta^K X^L f_{IJ}^T f_{KL}^Q g_{TQ} \\ &= \eta^I \eta^J \eta^K X^L f_{IJ}^T f_{TK}^Q g_{QL} \\ &= \frac{\lambda^2 + 2}{\lambda^2 + 1} \eta^I \eta^J \eta^K X^L f_{IT}^Q f_{JK}^T g_{QL} \\ &= \frac{\lambda^2 + 2}{\lambda^2 + 1} \eta^I \eta^J \eta^K X^L f_{TL}^Q f_{JK}^T g_{IQ} \\ &= \left(\frac{\lambda^2 + 2}{\lambda^2 + 1}\right)^2 \eta^I \eta^J \eta^K X^L f_{JT}^Q f_{KL}^T g_{IQ} \\ &= \left(\frac{\lambda^2 + 2}{\lambda^2 + 1}\right)^2 \eta^I \eta^J \eta^K X^L f_{IJ}^Q f_{KL}^T g_{QT}\end{aligned}$$

Thus,

$$\left\{1 - \left(\frac{\lambda^2 + 2}{\lambda^2 + 1}\right)^2\right\} \eta^I \eta^J \eta^K X^L f_{IJ}^T f_{KL}^Q g_{TQ} = 0$$

The first relation in (6.32) is now proved. The proof for the next relation can be performed analogously. Q.E.D.

Let Z^J be a field in $\mathcal{B}$ with the index n_z , and let X^J denote the field η^J or A^J with the index $n_x = -1$ or 1 , respectively. From Propositions 4 and 5 we obtain that if $n_z < n_x$:

$$\begin{aligned}\langle X, X, Z \rangle &= \langle Z, X, X \rangle \\ (-1)^{n_x n_z} \langle X, Z, X \rangle &= -(\lambda^2 + 1) \langle Z, X, X \rangle \\ \langle X, X, Z \rangle - (-1)^{n_x n_z} \langle X, Z, X \rangle &+ \langle Z, X, X \rangle \\ &= [3] \langle Z, X, X \rangle\end{aligned}\quad (6.34)$$

If $n_z > n_x$, we have:

$$\begin{aligned}
\langle Z, X, X \rangle &= ig\lambda\xi(\lambda^2+2) X^0\langle X, Z \rangle + (\lambda^2+1) \langle X, X, Z \rangle \\
(-1)^{n_x n_z} \langle X, Z, X \rangle &= - ig\lambda\xi X^0\langle X, Z \rangle - \langle X, X, Z \rangle \\
&= - (\lambda^2+2)^{-1} \{ \langle X, X, Z \rangle + \langle Z, X, X \rangle \} \\
\langle X, X, Z \rangle - (-1)^{n_x n_z} \langle X, Z, X \rangle + \langle Z, X, X \rangle & \\
&= - (-1)^{n_x n_z} [3] \langle X, Z, X \rangle
\end{aligned} \tag{6.35}$$

7. q -Deformed Chern-Simons and Cocycle Hierarchy

In the classical case Zumino [Zum1] [MSZ] introduced a homotopy operator k to compute the Chern-Simons. Generalizing his method we compute the second q -deformed Chern-Simons for $SU_q(2)$ in our previous paper [HHM]. Now, we compute the m -th q -deformed Chern-Simons for $SU_q(N)$.

Introduce a q -deformed homotopy operator k that is nilpotent and satisfies the Leibniz rule in the graded sense for the index n :

$$k^2 = 0, \quad d k + k d = \mathbf{1} \tag{7.1}$$

In the following we are going to show the existence of k , and compute the q -deformed Chern-Simons $Q_{2m-1}(A)$ from the m -th q -deformed Chern class by the operator k :

$$\begin{aligned}
P_m &= (d k + k d) P_m = d(k P_m) = d Q_{2m-1}(A) \\
Q_{2m-1}(A) &= k P_m
\end{aligned} \tag{7.2}$$

where we used $dP_m = 0$.

Introduce a real parameter t , $0 \leq t \leq 1$. When t changes from 0 to 1, the gauge potentials A_t^J change from 0 to A^J :

$$\begin{aligned}
A_t^J &= t A^J \\
F_t^J &= t dA^J + \{ig t^2/\lambda\} (A^0 A^J + A^J A^0) \\
&= t F^J + \{ig (t^2 - t)/\lambda\} (A^0 A^J + A^J A^0)
\end{aligned} \tag{7.3}$$

We choose the symmetrized definition for the q -deformed derivative and the q -deformed integral [GR] [Jackson] to fit our definition (2.9) for q -number. Define the

q -deformed derivative along t by:

$$\frac{\partial}{\partial_q t} f(t) = \frac{f(qt) - f(q^{-1}t)}{t(q - q^{-1})} \quad (7.4)$$

satisfying the q -deformed Leibniz rule:

$$\frac{\partial}{\partial_q t} f(t)g(t) = \frac{\partial f(t)}{\partial_q t} g(qt) + f(q^{-1}t) \frac{\partial g(t)}{\partial_q t} \quad (7.5)$$

The q -deformed integral is defined by:

$$\int_0^{t_0} d_q t f(t) = t_0(1 - q^2) \sum_{k=0}^{\infty} q^{2k} f(q^{2k+1}t_0) \quad (7.6)$$

At least for a polynomial, the q -deformed integral is the inverse of q -deformed derivative. For example,

$$\frac{\partial}{\partial_q t} t^m = [m] t^{m-1}, \quad \int_0^{t_0} d_q t t^{m-1} = t_0^m / [m]$$

Now, define the q -deformed Lie derivative $\hat{\delta}_q$ along t in the gauge space:

$$\hat{\delta}_q \equiv d_q t \frac{\partial}{\partial_q t} \quad (7.7)$$

and the q -deformed operator ℓ_t that satisfies the q -deformed Leibniz rule in the graded sense for the index n :

$$\begin{aligned} \ell_t A_t^J &= 0, \quad \ell_t F_t^J = \hat{\delta}_q A_t^J = d_q t A^J \\ \ell_t \{X_t^J Y_t^K\} &= \{\ell_t X_t^J\} Y_{qt}^K + (-1)^n X_{q^{-1}t}^J \{\ell_t Y_t^K\} \end{aligned} \quad (7.8)$$

where X_t^J and Y_t^K are the fields in $\mathcal{B}$, and X_t^J has the index n .

It is easy to check that for all formal polynomials (vanishing at $F_t^J = 0$ and $A_t^J = 0$) we have

$$\begin{aligned} \ell_t \ell_t &= 0 \\ \ell_t d + d \ell_t &= \hat{\delta}_q = d_q t \frac{\partial}{\partial_q t} \\ \hat{\delta}_q d &= d \hat{\delta}_q, \quad \hat{\delta}_q \ell_t = \ell_t \hat{\delta}_q \end{aligned} \quad (7.9)$$

Comparing it with (7.1) we obtain:

$$k = \int_0^1 \ell_t \quad (7.10)$$

The $(2m - 1)$ -th q -deformed Chern-Simons can be computed from (7.2) straightforwardly. In the following we give some examples. For $m = 2$ we have:

$$\begin{aligned}
\ell_t (P_2)_t &= \langle \ell_t F_t, F_{qt} \rangle + \langle F_{q^{-1}t}, \ell_t F_t \rangle \\
&= d_{qt} \{ \langle A, F_{qt} \rangle + \langle F_{q^{-1}t}, A \rangle \} \\
&= d_{qt} \{ t[2] \langle A, dA \rangle + ig\lambda^{-1}t^2(\lambda^2 + 2) \langle A, (A^0 A + A A^0) \rangle \} \\
&= d_{qt} \{ t[2] \langle A, dA \rangle + t^2 \xi^{-1}(\lambda^2 + 2) \langle A, A, A \rangle \}
\end{aligned} \tag{7.11}$$

where we have used (6.28).

$$\begin{aligned}
Q_3(A) &= k (P_2)_t \\
&= \langle A, dA \rangle + \xi^{-1} \{ [4]/[3][2] \} \langle A, A, A \rangle \\
&= \langle A, F \rangle - \{ \xi[3] \}^{-1} \langle A, A, A \rangle
\end{aligned} \tag{7.12}$$

For $m = 3$ we have:

$$\begin{aligned}
\ell_t (P_3)_t &= \langle \ell_t F_t, F_{qt}, F_{qt} \rangle + \langle F_{q^{-1}t}, \ell_t F_t, F_{qt} \rangle + \langle F_{q^{-1}t}, F_{q^{-1}t}, \ell_t F_t \rangle \\
&= d_{qt} \langle A, F_{qt}, F_{qt} \rangle + \langle F_{q^{-1}t}, A, F_{qt} \rangle + \langle F_{q^{-1}t}, F_{q^{-1}t}, A \rangle \\
&= d_{qt} t^2 \{ q^2 \langle A, dA, dA \rangle + \langle dA, A, dA \rangle + q^{-2} \langle dA, dA, A \rangle \} \\
&\quad + d_{qt} t^3 \xi^{-1} \{ (q^3 + q^{-1}) \langle A, A, A, dA \rangle + q^{-3} \langle A, A, dA, A \rangle \\
&\quad + q^3 \langle A, dA, A, A \rangle + (q + q^{-3}) \langle dA, A, A, A \rangle \}
\end{aligned}$$

$$\begin{aligned}
Q_5(A) &= k (P_3)_t \\
&= [3]^{-1} \{ q^2 \langle A, dA, dA \rangle + \langle dA, A, dA \rangle + q^{-2} \langle dA, dA, A \rangle \} \\
&\quad + \{ \xi[4] \}^{-1} \{ (q^3 + q^{-1}) \langle A, A, A, dA \rangle + q^{-3} \langle A, A, dA, A \rangle \\
&\quad + q^3 \langle A, dA, A, A \rangle + (q + q^{-3}) \langle dA, A, A, A \rangle \} \\
&= [3]^{-1} \{ q^2 \langle A, F, F \rangle + \langle F, A, F \rangle + q^{-2} \langle F, F, A \rangle \} \\
&\quad - ig \{ \xi[4][3] \}^{-1} \{ (q^3 + q^{-1}) \langle A, A, A, F \rangle + q \langle A, A, F, A \rangle \\
&\quad + q^{-1} \langle A, F, A, A \rangle + (q + q^{-3}) \langle F, A, A, A \rangle \}
\end{aligned} \tag{7.13}$$

Just like those in the classical case [HZ], the gauge fields F^J are invariant under the transformation:

$$A^J \rightarrow A^J - \eta^J, \quad d \rightarrow d + \delta \tag{7.14}$$

In fact,

$$\begin{aligned}
F^J &\rightarrow \mathcal{F}^J \\
&= (d + \delta) (A^J - \eta^J) \\
&\quad + \frac{ig}{\lambda} \{ (A^0 - \eta^0)(A^J - \eta^J) + (A^J - \eta^J)(A^0 - \eta^0) \} \\
&= F^J + \left\{ \delta A^J - d\eta^J - \frac{ig}{\lambda} (\eta^0 A^J + A^J \eta^0) \right\} \\
&\quad - \left\{ \delta \eta^J - \frac{ig}{\lambda} (\eta^0 \eta^J + \eta^J \eta^0 - A^0 \eta^J - \eta^J A^0) \right\} \\
&= F^J
\end{aligned}$$

Now, transforming (7.2) and expanding it by the ghost number, we obtain:

$$\begin{aligned}
P_m &= (d + \delta) Q_{2m-1}(A - \eta) \\
Q_{2m-1}(A - \eta) &= \sum_{n=0}^{2m-1} \omega_{2m-n-1}^n \\
P_m &= d\omega_{2m-1}^0 + \sum_{n=0}^{2m-2} \left\{ \delta \omega_{2m-n-1}^n + d\omega_{2m-n-2}^{n+1} \right\} + \delta \omega_0^{2m-1}
\end{aligned} \tag{7.15}$$

where the subscripts denote the degrees of form of the quantities, and the superscripts denote the ghost numbers. In two sides of (7.15) the quantities with the same degree of form and the same ghost number should be equal to each other, respectively:

$$\begin{aligned}
P_m &= d\omega_{2m-1}^0, \quad \delta \omega_0^{2m-1} = 0 \\
\delta \omega_{2m-n-1}^n + d\omega_{2m-n-2}^{n+1} &= 0, \quad n = 0, 1, \dots, (2m-2)
\end{aligned} \tag{7.16}$$

For example, for $m = 2$ we have:

$$Q_3(A - \eta) = \langle A - \eta, F \rangle - \{\xi[3]\}^{-1} \langle A - \eta, A - \eta, A - \eta \rangle$$

Simplifying them by the formulas given in Section 6, we obtain:

$$\begin{aligned}
\omega_3^0 &= Q_3(A) = \langle A, F \rangle - \{\xi[3]\}^{-1} \langle A, A, A \rangle \\
&= \langle A, dA \rangle + \frac{[4]}{\xi[3][2]} \langle A, A, A \rangle \\
\omega_2^1 &= - \langle \eta, dA \rangle - \{ig/\lambda\} \langle \eta, (A^0 A + A A^0) \rangle \\
&\quad + \{\xi[3]\}^{-1} \{ \langle \eta, A, A \rangle + \langle A, \eta, A \rangle + \langle A, A, \eta \rangle \} \\
&= - \langle \eta, dA \rangle \\
\omega_1^2 &= - \{\xi[3]\}^{-1} \{ \langle \eta, \eta, A \rangle + \langle \eta, A, \eta \rangle + \langle A, \eta, \eta \rangle \} \\
&= - \xi^{-1} \langle \eta, A, \eta \rangle \\
\omega_0^3 &= \{\xi[3]\}^{-1} \langle \eta, \eta, \eta \rangle
\end{aligned} \tag{7.17}$$

It is easy to check by the formulas in Section 6 that (7.17) satisfies (7.16).

8. q -deformed Lagrangian and Yang-Mills equation

In the present paper the spacetime is the ordinary commutative Minkowski spacetime. Explicitly writing down the spacetime indices, we have:

$$A^J = A_\mu^J dx^\mu, \quad F^J = \frac{1}{2} F_{\mu\nu}^J dx^\mu \wedge dx^\nu \quad (8.1)$$

It is well known that the metric $g^{\mu\nu}$ in the Minkowski spacetime can change the covariant index to contravariant index, or vice versa.

Now, the q -deformed Lagrangian that is covariant both in the Lorentz transformation and the q -gauge transformation is:

$$\mathcal{L} = -\frac{1}{4} \langle F_{\mu\nu}, F^{\mu\nu} \rangle = -\frac{1}{4} (F^I)_{\mu\nu} (F^J)^{\mu\nu} g_{IJ} \quad (8.2)$$

We have known that the components of the identity and the adjoint representations are separated in the q -deformed Chern class, and obviously in the q -deformed Lagrangian. Here we only discuss the q -deformed Lagrangian constructed by the adjoint components.

The q -deformed Yang-Mills equation is just the q -deformed Lagrangian equation:

$$\begin{aligned} & \partial_\mu (F^J)^{\mu\nu} (g_{KJ} + g_{JK}) \\ &= (\lambda^2 + 2)^{-1} \left(\mathbf{C}_{RK}^I - \mathbf{C}_{KR}^I \right) \left\{ g_{IJ} (A^R)_\mu (F^J)^{\mu\nu} + g_{JI} (F^J)^{\mu\nu} (A^R)_\mu \right\} \end{aligned} \quad (8.3)$$

Acknowledgments. This work was supported by the National Natural Science Foundation of China and Grant No. LWTZ-1298 of Chinese Academy of Sciences.

References

- [AC] P. Aschieri, L. Castellani, An introduction to noncommutative differential geometry on quantum groups, *Inter. J. Mod. Phys.* **A8**(1993)1667-1706.

- [AV] I. Ya. Aref'eva and I. V. Volovich, Quantum group chiral fields and differential Yang-Baxter equations, *Phys. Lett.* **B264**(1991)62-68.
- [Bax] R. J. Baxter, Partition function of the eight-vertex lattice model, *Ann. Phys.* **70**(1972)193-228.
- [Ber] D. Bernard, Quantum Lie Algebras and Differential Calculus on Quantum Groups – Proc. 1990 Yukawa Int. Seminar (kyoto); *Phys. Lett.* **B260**(1991)389.
- [Cas] L. Castellani, $U_q(N)$ Gauge Theories, Preprint, DFTT-74/92, 1992.
- [Con] Alain Connes, Non-commutative differential geometry, *Publ. Math. IHES.* **62**(1986)41-144.
- [CSWW] U. Carow-Watamura, M. Schlieker, S. Watamura and W. Weich, Bicovariant differential calculus on quantum groups $SU_q(N)$ and $SO_q(N)$, *Commun. Math. Phys.* **142**(1991)605-641.
- [CW] U. Carow-Watamura and S. Watamura, Complex quantum group, dual algebra and bicovariant differential calculus, *Commun. Math. Phys.* **151**(1993)487-514.
- [FP] L. D. Faddeev, P. N. Pyatov, The differential calculus on quantum linear groups, hep-th/9402070.
- [FRT] L. D. Faddeev, N. Y. Reshetikhin and L. A. Takhtajan, Quantization of Lie groups and Lie algebras, in *Algebraic analysis*, Academic Press, 129, 1988.
- [GJ] D. Gross and R. Jackiw, Effect of anomalies on quasi-renormalizable theories, *Phys. Rev.* **D6**(1972)477-493.
- [GR] G. Gasper and M. Rahman, Basic Hypergeometric Series, in *Encyclopedia of Mathematics and its Applications 35*, Cambridge, 1990.
- [HHM] Bo-Yu Hou, Bo-Yuan Hou and Zhong-Qi Ma, q -deformed Chern class, Chern-Simons and Cocycle hierarchy, preprint BIHEP-TH-94-9, hep-th/9403153.
- [HZ] Bo-Yu Hou and Yao-Zhong Zhang, Cohomology in connection space, family index theorem, and abelian gauge structure, *J. Math. Phys.* **28**(1987)1709-1715.
- [Isa] A. P. Isaev, Quantum group covariant noncommutative geometry, Dubna Preprint JINR E2-94-38, hep-th/9402060.

- [IP] A. P. Isaev and Z. Popowicz, q -trace for the quantum groups and q -deformed Yang-Mills theory, Wroclaw preprint ITP UWr 786/91.
- [Jackiw] R. Jackiw, New directions for topological research on gauge theories, in *Symposium on Anomalies, Geometry, Topology*, P.156, Eds. W. A. Bardeen and A. R. White, World Scientific, Singapore, 1985.
- [Jackson] F. H. Jackson, *Quart. J. Pure Appl. Math.* **41**(1910)193.
- [Jur] B. Jurčo, Differential calculus on quantized simple Lie groups, *Lett. Math. Phys.* **22**(1991)177-186.
- [KR] A. N. Kirillov and N. Y. Reshetikhin, q -Weyl group and a multiplicative formula for universal R -matrices, HUTMP 90/B261.
- [Ma] Zhong-Qi Ma, *Yang-Baxter Equation and Quantum Enveloping algebras*, World Scientific, Singapore, 1993.
- [Manin] Yu. I. Manin, Quantum Groups and Non-Commutative Geometry, Montreal Univ. preprint, CRM-1561, 1988.
- [MSZ] J. Mañes, R. Stora and B. Zumino, Algebraic study of chiral anomaly, *Commun. Math. Phys.* **102**(1985)157-174.
- [Res] N. Y. Reshetikhin, Quantum universal enveloping algebras, the Yang-Baxter equation and invariant of links I, Preprint, LOMI, E-4-87, 1987.
- [SW] X. D. Sun and S. K. Wang, Bicovariant differential calculus on quantum group $GL_q(n)$, Worldlab-Beijing preprint CCAST-92-04, 1992.
- [SWZ] P. Schupp, P. Watts and B. Zumino, Differential geometry on linear quantum groups, preprint, LBL-32314, UCB-PTH-92/13.
- [Wat] S. Watamura, Quantum deformation of BRST algebra, *Commun. Math. Phys.* **158**(1993)67-92.
- [Wor1] S. L. Woronowicz, Compact matrix pseudogroups, *Commun. Math. Phys.* **111**(1987)613-665.
- [Wor2] S. L. Woronowicz, Differential calculus on compact matrix pseudogroups (quantum groups), *Commun. Math. Phys.* **122**(1989)125-170.
- [WZ] J. Wess and B. Zumino, Covariant differential calculus on the quantum hyperplane, *Nucl. Phys. (Proc. Suppl.)* **B18**(1990)302-312.

- [Yang] C. N. Yang, Some exact results for the many-body problem in one dimension with repulsive delta-function, *Phys. Rev. Lett.* **19**(1967)1312-1315.
- [Zum1] B. Zumino, Chiral Anomalies and Differential Geometry, Lectures given at Les Houches, Preprint, LBL-16747, 1983.
- [Zum2] B. Zumino, Introduction to the differential geometry of quantum groups, LBL-31432 and UCB-PTH-62-91, Notes of a plenary talk given at the 10th IAMP Conf., Leipzig, 1991.